

Quasi-hereditary algebras via generator-cogenerators of local self-injective algebras and transfer of Ringel duality

Daiva Pučinskaitė

Abstract

The dominant dimension of algebras in the class $\mathbf{A}$ of 1-quasi-hereditary algebras introduced in [9] is at least two. By the Morita-Tachikawa Theorem this implies that $\mathbf{A}$ is related to a certain class $\mathbf{B}$ of algebras via bimodules satisfying the double centralizer condition. In this paper we specify the class $\mathbf{B}$ and the modules over algebras in $\mathbf{B}$ connected with $\mathbf{A}$.

The class $\mathbf{A}$ is not closed under taking the Ringel-dual. However the dominant dimension of the Ringel-dual $R(A)$ of $A \in \mathbf{A}$ is at least two. This fact induces a corresponding concept of modules over $B \in \mathbf{B}$ which yield the algebras A and $R(A)$ for $A \in \mathbf{A}$.

Introduction

Let $\mathcal{A}, \mathcal{B}$ be algebras. An $\mathcal{A}\mathcal{B}$ -bimodule ${}_{\mathcal{A}}\mathcal{M}_{\mathcal{B}}$ satisfying the double centralizer condition $\mathcal{A} \cong \text{End}_{\mathcal{B}}(\mathcal{M}_{\mathcal{B}})$ and $\mathcal{B} \cong \text{End}_{\mathcal{A}}({}_{\mathcal{A}}\mathcal{M})$ provides a relationship between the representation theories of the algebras $\mathcal{A}$ and $\mathcal{B}$ that may differ in terms of their homological properties. Soergel's 'Struktursatz' relating an algebra $\mathcal{A}_{\Theta}(\mathfrak{g})$ corresponding to a block Θ of the Bernstein-Gelfand-Gelfand category $\mathcal{O}(\mathfrak{g})$ of a complex semisimple Lie algebra $\mathfrak{g}$ with a subalgebra of the corresponding coinvariant algebra as well as the Schur-Weyl duality between the Schur algebra $S(n, r)$ for $n \geq r$ and the group algebra $K\Sigma_r$ of the symmetric group are prominent examples for this connection (see for instance [7]).

In this paper we present a further example for this situation which relates a 1-quasi-hereditary algebra A defined in [9] with a local self-injective algebra B via an A - B -bimodule L whose structure has a precise description: The A -module L is a projective-injective indecomposable and the $\text{End}_A(L)$ -module L decomposes into a direct sum of local ideals of $B := \text{End}_A(L)$ generated by the endomorphisms corresponding to certain paths in the quiver of A (see Proposition 2.1.1). The B -module L is a generator-cogenerator of $\text{mod } B$ (i.e., any projective resp. injective indecomposable B -module is a direct summand of L). Thus, any 1-quasi-hereditary algebra can be defined as an endomorphism algebra of a generator-cogenerator of a local self-injective algebra.

The algebras $\mathcal{A}_{\Theta}(\mathfrak{g})$ and $S(n, r)$ belong to the class $\mathscr{A}$ of quasi-hereditary algebras with a duality (induced by an anti-automorphism) and with dominant dimension at least two (see [7] and [8]). The class of 1-quasi-hereditary algebras has a non-empty intersection with $\mathscr{A}$. Many factor algebras of $\mathcal{A}_{\Theta}(\mathfrak{g})$ related to certain saturated subsets of weights are 1-quasi-hereditary. Note that a 1-quasi-hereditary algebra does not have a duality in general. The results in this paper clarify the connection between $\mathcal{A}_{\Theta}(\mathfrak{g})$ and the coinvariant algebras.

Our first result presents a one-to-one correspondence (up to isomorphism) between the class of 1-quasi-hereditary algebras (over an algebraically closed field K) and the class of pairs (B, L) yielding 1-quasi-hereditary algebras via the double centralizer condition, where B is a local self-injective algebra and $L \in \text{mod } B$ satisfies certain properties.

Partly supported by the D.F.G. priority program SPP 1388 "Darstellungstheorie".

Any algebra $\mathcal{A}$ in this paper is basic, thus given by a quiver and relations $(Q(\mathcal{A}), \mathcal{I}(\mathcal{A}))$.

Theorem A. *Let A, B be finite dimensional basic K -algebras and L be a B -module. Let $n \in \mathbb{N}$ and $(\Lambda := \{1, \dots, n\}, \leq)$ be a partially ordered set. The following statements are equivalent:*

- (i) *A with $(\Lambda, \leq)$ is 1-quasi-hereditary (here we identify Λ with the vertices in $Q(A)$), i.e., $A \cong \text{End}_B(L)^{op}$, where L is a multiplicity-free generator-cogenerator of $\text{mod } B$.*
- (ii) *B is local, self-injective with $\dim_K B = n$ and $L \cong \bigoplus_{i \in \Lambda} L(i)$ where $L(i)$ are local submodules of B and $L(1) = B$, moreover, for all $i, j \in \Lambda$ the following properties hold:*
 - (a) *$L(i) \twoheadrightarrow L(j)$ if and only if $i \leq j$,*
 - (b) *$\text{rad}(L(i)) = \sum_{i < j} L(j)$.*

An algebra of the form $A_\Theta(\mathfrak{g})$ is 1-quasi-hereditary if $\text{rank}(\mathfrak{g}) \leq 2$, hence Theorem A is applicable for these algebras.

Dlab, Heath and Marko have shown in [4] that a pair (B, L) with the properties in (ii) yields a quasi-hereditary BGG-algebra (defined by Irving in [6]) if B is commutative. The next theorem strengthens the main theorem in [4] by determining the properties of a 1-quasi-hereditary algebra $A \cong \text{End}_B(L)^{op}$ for which B is commutative.

In the quiver Q of a 1-quasi-hereditary algebra A between two vertices i and j either there are no arrows or $i \leq j$ (see [9, Theorem 2.7]). Thus for any path p in Q the opposite path p^{op} also belongs to Q .

Theorem B. *Let $A \cong \text{End}_B(L)^{op}$ be a 1-quasi-hereditary algebra where the B -module L satisfies the conditions (ii) in Theorem A. The following statements are equivalent:*

- (i) *B is commutative.*
- (ii) *If ρ is a relation of A , then ρ^{op} is also a relation of A .*
- (iii) *A has a duality induced by the anti-automorphism $p \mapsto p^{op}$.*

The coinvariant algebra related to the algebra $\mathcal{A}_\Theta(\mathfrak{g})$ is commutative. Note that Theorem B is also true for all algebras $\mathcal{A}_\Theta(\mathfrak{g})$, where the quiver and relations are known, also for non 1-quasi-hereditary algebras (see [13]).

The concept of Ringel duality introduced in [11] is essential in the theory of quasi-hereditary algebras (for a (basic) quasi-hereditary algebra $\mathcal{A}$ there exists another quasi-hereditary algebra $R(\mathcal{A})$ such that $R(R(\mathcal{A})) \cong \mathcal{A}$). In this paper we show how the Ringel duality $R(-)$ induces a corresponding concept $\tilde{R}(-)$ on certain generator-cogenerators of a local self-injective algebra. This is based on the fact that the class of 1-quasi-hereditary algebras is not closed under Ringel duality, however for any 1-quasi-hereditary algebra A there exists an $R(A)$ - $\tilde{R}(B)$ -bimodule $\tilde{R}(L)$ having the double centralizer condition (see Lemma 4.1).

Our next result explicitly determines the aforementioned correspondence for those 1-quasi-hereditary algebras, whose Ringel duals are also 1-quasi-hereditary.

Theorem C. *Let $(A, \leq)$ and $(R(A), \geq)$ be 1-quasi-hereditary algebras as well as (B, L) and $(\tilde{R}(B), \tilde{R}(L))$ the corresponding pairs (w.r.t. Theorem A (ii)), where $L \cong \bigoplus_{i \in \Lambda} L(i)$ and $\tilde{R}(L) \cong \bigoplus_{i \in \Lambda} \tilde{R}(L(i))$. Then $B \cong \tilde{R}(B)$ and for every $i \in \Lambda$ we have*

$$\tilde{R}(L(i)) \cong B / \left(\sum_{j \not\leq i} L(j) \right) \cong \bigcap_{j \not\leq i} \ker (B \twoheadrightarrow L(j)).$$

In particular, if a 1-quasi-hereditary algebra is Ringel self-dual, then $L \cong \tilde{R}(L)$. There exists a permutation $\sigma \in \text{Sym}(\dim_K B)$ with $L(\sigma(i)) \cong \tilde{R}(L(i))$. The algebras of the form $\mathcal{A}_\Theta(\mathfrak{g})$ are Ringel self-dual. In case of $\text{rank}(\mathfrak{g}) \leq 2$ we have this situation.

The paper is organized as follows: In Section 1, we introduce the Morita-Tachikawa Theorem which shows that a minimal faithful module over an algebra of dominant dimension at least two has the double centralizer property. The results of this paper rest on this theorem. We also recall the relevant definitions and give some examples which show the diversity of modules over a local self-injective algebra satisfying the double centralizer condition.

Section 2 is devoted to the proof of Theorem A. The paths in the quiver of a 1-quasi-hereditary algebra of the form $p(j, i, k)$ defined in [9, Section 3] play an important role. The other part of the proof is based on the structure of the B -module L which will be analyzed in Lemma 2.2.1. We also determine an easier transition from the B -maps of L to the relations of the algebra $A = \text{End}_B(L)^{op}$ (see Remark 2.2.5). This is used in the proof of Theorem B in Section 3.

In Section 4 we describe the transfer of Ringel duality. Consequently we obtain a decomposition of the class of 1-quasi-hereditary algebras with their Ringel duals into subclasses which are closed under Ringel duality. Moreover, the algebras in a fixed subclass arise from the same local self-injective algebra. Subsequently we present the proof of Theorem C.

1. Preliminaries

Unless otherwise specified, any algebra $\mathcal{A}$ is an associative, finite dimensional, basic K -algebra over an algebraically closed field K . Thus $\mathcal{A}$ is determined by a quiver Q and relations $\mathcal{I}$, i.e., $\mathcal{A} = KQ/\mathcal{I}$ (Theorem of Gabriel). Furthermore, $\text{mod } \mathcal{A}$ is the category of finite dimensional left $\mathcal{A}$ -modules. For $\mathcal{M} \in \text{mod } \mathcal{A}$ we denote by $\text{soc } \mathcal{M}$, $\text{rad } \mathcal{M}$ and $\text{top } \mathcal{M}$ the socle, the radical and the top of $\mathcal{M}$, respectively, and $\text{add}(\mathcal{M})$ is the full subcategory of $\text{mod } \mathcal{A}$ whose objects are the direct sums of direct summands of $\mathcal{M}$ (for references see [1] or [2]). To distinguish between an arbitrary algebra $\mathcal{A}$ and a 1-quasi-hereditary algebra, we denote the latter by A .

We repeat some notations and facts about bound quiver algebras $\mathcal{A} = KQ/\mathcal{I}$. Throughout, we identify the set of vertices $Q_0 := Q_0(\mathcal{A})$ with the set $\Lambda = \{1, \dots, |Q_0|\}$. The product of arrows $(k \rightarrow i)$ and $(i \rightarrow j)$ is given by $(k \rightarrow i \rightarrow j) = (i \rightarrow j) \cdot (k \rightarrow i)$. We denote by $P(i)$, $I(i)$, $S(i)$ and e_i the projective indecomposable, injective indecomposable, simple $\mathcal{A}$ -module and the primitive idempotent, respectively, corresponding to $i \in \Lambda$. Let $\mathcal{M} \in \text{mod } \mathcal{A}$, then for each vertex i there exists a K -subspace of $\mathcal{M}$ corresponding to i , denoted by $\mathcal{M}_i$. We have $\mathcal{M}_i \cong \text{Hom}_{\mathcal{A}}(P(i), \mathcal{M})$ via $m \mapsto f_{(m)} : P(i) \rightarrow \mathcal{M}$, where

$f_{(m)}(a \cdot e_i) = a \cdot m$ for all $a \in \mathcal{A}$. Recall that $\text{End}_{\mathcal{A}}(\mathcal{M})^{op}$ is a K -algebra with product $F \circ G = \left(\mathcal{M} \xrightarrow{G} \mathcal{M} \xrightarrow{F} \mathcal{M} \right)$. The (left) $\text{End}_{\mathcal{A}}(\mathcal{M})^{op}$ -module $\mathcal{M}$ (written ${}_{\text{End}_{\mathcal{A}}(\mathcal{M})^{op}}\mathcal{M}$) is isomorphic to $\bigoplus_{i \in \Lambda} \text{Hom}_{\mathcal{A}}(P(i), \mathcal{M})$. The Jordan-Hölder multiplicity of $S(i)$ in $\mathcal{M}$ is denoted by $[\mathcal{M} : S(i)]$. In particular, we have $\dim_K \text{Hom}_{\mathcal{A}}(P(i), \mathcal{M}) = [\mathcal{M} : S(i)]$.

1.1. Theorem of Morita-Tachikawa

Based on various works by Morita and Tachikawa, Ringel has described in [12] a relationship between the algebras $\mathcal{A}$ and $\text{End}_{\mathcal{A}}(\mathcal{M})^{op}$ via an $\mathcal{A}$ -module $\mathcal{M}$ having the double centralizer condition. We recall some notations and terminology needed in the statement of the theorem below. Let $\mathcal{B}$ be an artin (basic) algebra, then the *dominant dimension* of $\mathcal{B}$ is at least 2 (written $\text{dom. dim } \mathcal{B} \geq 2$), if there exists an exact sequence $0 \rightarrow {}_{\mathcal{B}}\mathcal{B} \rightarrow \mathcal{M}_1 \rightarrow \mathcal{M}_2$ such that $\mathcal{M}_1, \mathcal{M}_2$ are projective and injective $\mathcal{B}$ -modules (all finite dimensional algebras are artin algebras). A $\mathcal{B}$ -module $\mathcal{N}$ is called *minimal faithful* if $\mathcal{N}$ is faithful (i.e., ${}_{\mathcal{B}}\mathcal{B}$ can be embedded into a direct sum of copies of $\mathcal{N}$) and $\mathcal{N}$ is a direct summand of any faithful $\mathcal{B}$ -module. A minimal faithful $\mathcal{B}$ -module is unique (up to isomorphism) and will be denoted by $\mathcal{F}(\mathcal{B})$. A $\mathcal{B}$ -module $\mathcal{M}$ is a *generator-cogenerator* of $\text{mod } \mathcal{B}$ if every projective indecomposable as well as every injective indecomposable $\mathcal{B}$ -module is a direct summand of $\mathcal{M}$. We denoted by $[\mathcal{B}]$ resp. $[\mathcal{B}, \mathcal{M}]$ the isomorphism class of $\mathcal{B}$ and a $\mathcal{B}$ -module $\mathcal{M}$.

1.1 Theorem (Morita-Tachikawa). *There is are bijections Ψ and Φ between*

$$\begin{aligned} \mathbf{X} &:= \left\{ [\mathcal{A}] \mid \mathcal{A} \text{ is a basic artin algebra, } \text{dom. dim } \mathcal{A} \geq 2 \right\} \text{ and} \\ \mathbf{Y} &:= \left\{ [\mathcal{B}, \mathcal{M}] \mid \begin{array}{l} \mathcal{B} \text{ is a basic artin algebra,} \\ \mathcal{M} \text{ is a multiplicity-free, generator-cogenerator of } \text{mod } \mathcal{B} \end{array} \right\} \end{aligned}$$

defined as follows:

$$\begin{array}{ccc} \mathbf{X} & \xrightarrow{\Phi} & \mathbf{Y} \\ [\mathcal{A}] & \mapsto & [\mathcal{B}(\mathcal{A}) := \text{End}_{\mathcal{A}}(\mathcal{F}(\mathcal{A}))^{op}, {}_{\mathcal{B}(\mathcal{A})}\mathcal{F}(\mathcal{A})] \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathbf{Y} & \xrightarrow{\Psi} & \mathbf{X} \\ [\mathcal{B}, \mathcal{M}] & \mapsto & [\text{End}_{\mathcal{B}}(\mathcal{M})^{op}] \end{array},$$

such that $\Psi \circ \Phi = \text{Id}_{\mathbf{X}}$ and $\Phi \circ \Psi = \text{Id}_{\mathbf{Y}}$.

This theorem also provides correspondences between the subsets of $\mathbf{X}$ and their image under Φ in $\mathbf{Y}$. By the Theorem of König, Slungård and Xi [7, Theorem 1.3] the set of isomorphism classes of algebras which corresponds to the blocks of the BGG-category $\mathcal{O}$ is a subset of $\mathbf{X}$. Theorem 1.1 restricted to this subset is known as Soergel's 'Struktursatz'. Moreover, the dominant dimension of Schur algebras $S(n, r)$ (with $n > r$) is at least two. The Schur-Weyl duality is a special case of Theorem 1.1. The module $\mathcal{F}(\mathcal{A})$ has the double centralizer condition.

Note that the Theorem of Morita-Tachikawa provides some connections between a finite dimensional algebra $\mathcal{A} = KQ/\mathcal{I}$ with $\text{dom. dim } \mathcal{A} \geq 2$ and a pair $(\mathcal{B}, \mathcal{M})$ with $\Phi[A] = [\mathcal{B}, \mathcal{M}]$: The $\mathcal{B}$ -module $\mathcal{M}$ has $|Q_0|$ pairwise non-isomorphic, indecomposable direct summands (they correspond to the vertices of Q). Because $\mathcal{A} \hookrightarrow \mathcal{F}(\mathcal{A})^m$ for some $m \in \mathbb{N}$, we have $[\mathcal{F}(\mathcal{A}) : S(i)] \neq 0$ for all $i \in Q_0$, thus $\text{Hom}_{\mathcal{A}}(P(i), \mathcal{F}(\mathcal{A})) \neq 0$. Since the $\mathcal{B}$ -module $\mathcal{M} \cong \mathcal{F}(\mathcal{A})$ is isomorphic to $\bigoplus_{i \in Q_0} \text{Hom}_{\mathcal{A}}(P(i), \mathcal{F}(\mathcal{A}))$, we obtain that the $\text{End}_{\mathcal{A}}(\mathcal{F}(\mathcal{A}))^{op}$ -module $\text{Hom}_{\mathcal{A}}(P(i), \mathcal{F}(\mathcal{A}))$ is indecomposable for any $i \in Q_0$.

1.2. Quasi-hereditary algebras and local self-injective algebras

In this paper we consider a subclass of quasi-hereditary algebras with dominant dimension at least 2 which are related to local self-injective algebras. We recall some necessary definitions.

Quasi-hereditary algebras were defined by Cline, Parshall and Scott in [3]. We use the equivalent definition and terminology given by Dlab and Ringel in [5]: Let $\mathcal{A}$ be a bound quiver algebra and $(\Lambda, \leq)$ a poset (with $\Lambda = Q_0(\mathcal{A})$). For every $i \in \Lambda$ the *standard* module $\Delta(i)$ is the largest factor module of $P(i)$ such that $[\Delta(i) : S(k)] = 0$ for all $k \in \Lambda$ with $k \not\leq i$. The modules in the full subcategory $\mathfrak{F}(\Delta)$ of $\text{mod } \mathcal{A}$ consisting of the modules having a filtration such that each subquotient is isomorphic to a standard module are called Δ -good and these filtrations are Δ -good filtrations. For $M \in \mathfrak{F}(\Delta)$ we denote by $(M : \Delta(i))$ the (well-defined) number of subquotients isomorphic to $\Delta(i)$ in a Δ -good filtration of M .

An algebra $\mathcal{A}$ with $(\Lambda, \leq)$ is *quasi-hereditary* if for all $i, k \in \Lambda$ the following conditions are satisfied:

- $[\Delta(i) : S(i)] = 1$,
- $P(i)$ is a Δ -good module with $(P(i) : \Delta(k)) = 0$ for all $k \not\leq i$ and $(P(i) : \Delta(i)) = 1$.

Throughout, $(\mathcal{A}, \leq)$ denotes an algebra $\mathcal{A}$ with a partial order $\leq$ on the vertices $\Lambda = Q_0(\mathcal{A})$.

We can identify the vertices of the quivers of $\mathcal{A}$ and $\mathcal{A}^{op}$. An algebra $(\mathcal{A}, \leq)$ is quasi-hereditary if and only if $(\mathcal{A}^{op}, \leq)$ is quasi-hereditary (see [5]). The standard duality $\mathcal{D} := \text{Hom}_K(-, K)$ provides the *costandard* $\mathcal{A}$ -module $\nabla(i) \cong \mathcal{D}(\Delta_{\mathcal{A}^{op}}(i))$ corresponding to $i \in \Lambda$ and also the subcategory $\mathfrak{F}(\nabla)$ of $\text{mod } \mathcal{A}$ of all ∇ -good modules.

1.2.1 Remark. Let $(\mathcal{A}, \leq)$ be a quasi-hereditary algebra and let M be a projective-injective $\mathcal{A}$ -module. Moreover, let $\text{soc}(\Delta(i)) \in \text{add}(\text{soc } M)$ and $P(i) \hookrightarrow M$ with $M/P(i) \in \mathfrak{F}(\Delta)$ for all $i \in \Lambda$. In this case we have $\text{soc}(M/P(i)) \in \text{add}\left(\bigoplus_{j \in \Lambda} \text{soc } \Delta(j)\right) \subseteq \text{add}(\text{soc } M)$ and consequently $M/P(i)$ can be embedded into some copies of M for any $i \in \Lambda$. In other words, there exist $m, r \in \mathbb{N}$ such that the sequence $0 \rightarrow \mathcal{A} \rightarrow M^m \rightarrow M^r$ is exact, i.e., we have $\text{dom. dim } \mathcal{A} \geq 2$.

We recall the definition of a 1-quasi-hereditary algebra from [9] and show that we have the situation described in the foregoing remark.

1.2.2 Definition (1-quasi-hereditary). A quasi-hereditary algebra A with $(\Lambda, \leq)$ is called *1-quasi-hereditary* if for all $i, j \in \Lambda = \{1, \dots, n\}$ the following conditions are satisfied:

- (1) There is a smallest and a largest element with respect to $\leq$,
without loss of generality we will assume them to be 1 resp. n ,
- (2) $[\Delta(i) : S(j)] = (P(j) : \Delta(i)) = 1$ for $j \leq i$,
- (3) $\text{soc } P(j) \cong \text{top } I(j) \cong S(1)$,
- (4) $\Delta(i) \hookrightarrow \Delta(n)$ and $\nabla(n) \twoheadrightarrow \nabla(i)$.

1.2.3 Lemma. Let $(A, \leq)$ be a 1-quasi-hereditary algebra with $\{1\} = \min(\Lambda, \leq)$. Then $P(1)$ is a minimal faithful A -module and $\text{dom. dim } A \geq 2$.

Proof. According to [9, 2.6] we have $P(1) \cong I(1)$ and $\Delta(j) \hookrightarrow P(i) \hookrightarrow P(1)$ for every $i, j \in \Lambda$. We obtain ${}_A A \hookrightarrow P(1)^{|\Lambda|}$, thus $P(1)$ is a minimal faithful A -module because $P(1)$ is indecomposable (we have $\mathcal{F}(A) \cong P(1)$).

Since $\text{soc } \Delta(i) \cong \text{soc } P(1) \cong S(1)$ and $P(1)/P(i) \in \mathfrak{F}(\Delta)$ (see [9, 4.3]), we have $\text{dom. dim } A \geq 2$ according to Remark 1.2.1. $\square$

An (finite dimensional, basic) algebra $\mathcal{B}$ is local and self-injective if and only if the socle and the top of $\mathcal{B}$ -module $\mathcal{B}$ are simple. An ideal I of $\mathcal{B}$ is a two-sided, local ideal if $\mathcal{B} \cdot I \subseteq I$ as well as $I \cdot \mathcal{B} \subseteq I$ and $\text{rad}(I)$ is the (uniquely determined) maximal submodule of I .


1.2.4 Definition ($\boxed{\leq}$). Let B be an algebra, $L \in \text{mod } B$ and $(\Lambda = \{1, \dots, n\}, \leq)$ be a poset. We say that the pair (B, L) satisfies the condition $\boxed{\leq}$ if

- (1) B is a local, self-injective algebra, $\dim_K B = n$,
- (2) $L \cong \bigoplus_{i \in \Lambda} L(i)$, where $L(1), \dots, L(n)$ are two-sided local ideals of B with $L(1) = B$ and for all $i, j \in \Lambda$ the following is satisfied:
 - (a) $L(i) \twoheadrightarrow L(j)$ if and only if $i \leq j$,
 - (b) $\text{rad}(L(i)) = \sum_{i < j} L(j)$.

1.2.5 Remark. If (B, L) satisfies the condition $\boxed{\leq}$, then $[B, L] \in \mathbf{Y}$: Since B is local and self-injective, any projective resp. injective, indecomposable B -module is isomorphic to ${}_B B$. Hence L is a generator-cogenerator of $\text{mod } B$, because $B (= L(1))$ is a direct summand of L . The condition (a) implies $L(i) \cong L(j)$ if and only if $i = j$, therefore L is multiplicity-free. Moreover, the quiver of B consists of one vertex and finitely many loops (B is isomorphic to $K \langle x_1, \dots, x_r \rangle / J$, where J is an ideal with $J \subseteq \langle x_1, \dots, x_r \rangle^2$ and $r \in \mathbb{N}$).

It should be noted that for a fixed local, self-injective algebra B a poset $(\Lambda, \leq)$ and a B -module L with the condition $\boxed{\leq}$ are not uniquely determined.

1.2.6 Example. The algebra $B = \mathbb{C}[x, y] / \langle xy, x^3 - y^3 \rangle$ is local and self-injective with $\dim_{\mathbb{C}} B = 6$. The next diagrams present the partial orders $\leq_{(i)}$ on $\Lambda = \{1, \dots, 6\}$ and the generators-cogenerators $L_i = \bigoplus_{k=1}^6 L_i(k)$ of $\text{mod } B$ for $i = 1, 2, 3$. We write $L_i(k) \rightarrow L_i(k')$ if $L_i(k) \supset L_i(k')$ and there does not exist a submodule U with $L_i(k) \supset U \supset L_i(k')$. In other words, $k <_{(i)} k'$ and k, k' are neighbours for all $1 \leq i \leq 3$ and $1 \leq k, k' \leq 6$. In the diagram on the right-hand side, $\mathbf{i} \in \mathbb{C}$ is such that $\mathbf{i}^2 = -1$ and $\omega = \frac{1}{2} + \mathbf{i} \frac{\sqrt{3}}{2}$ is a 6th root of unity.



It is easy to check that (B, L_i) satisfies the condition $\boxed{\leq_{(i)}}$ and therefore $[B, L_i] \in \mathbf{Y}$. In

view of Theorem A, L_i is an A_i - B -bimodule, where $A_i = \text{End}_B(L_i)^{op}$ is a 1-quasi-hereditary algebra for $i = 1, 2, 3$ (the quiver and relations of A_i are given in 3.2). Note that the algebra A_3 is associated to a regular block of the BGG-category $\mathcal{O}(\mathfrak{sl}_3)$.

In the next section we prove Theorem A which can be rewritten as follows:

Theorem A. *Let A, B be finite dimensional basic K -algebras and $n \in \mathbb{N}$. Moreover, let $(\Lambda := \{1, \dots, n\}, \leq)$ be partially ordered. The following statements are equivalent:*

(i) *The algebra $(A, \leq)$ is 1-quasi-hereditary, i.e., $A \cong \text{End}_B(L)^{op}$ and L is a multiplicity-free generator-cogenerator of $\text{mod } B$,*

(ii) *The pair (B, L) satisfies the condition $\boxed{\leq}$.*

For any 1-quasi-hereditary algebra A there exists an (up to isomorphism) uniquely determined algebra B and a multiplicity-free generator-cogenerator L of $\text{mod } B$ with $A \cong \text{End}_B(L)^{op}$ (see Theorem 1.1 and Lemma 1.2.3). According to Remark 1.2.5, Theorem A provides bijections between the isomorphism classes of 1-quasi-hereditary algebras and the pairs defined in 1.2.4.

1.3. BGG-algebras

Dlab, Heath and Marko have shown in [4] that if for a commutative algebra B and a B -module L the pair (B, L) satisfies the condition $\boxed{\leq}$, then the algebra $\text{End}_B(L)^{op}$ is a BGG-algebra as defined by Irving in [6]. Our Theorem B elaborates on the Theorem in [4].

We refer to the definition of BGG-algebras given by Xi in [14] (these algebras are also BGG-algebras in the sense of [6]): A quasi-hereditary algebra $\mathcal{A}$ is called a *BGG-algebra* if there is a duality δ of $\text{mod } \mathcal{A}$ such that δ induces a K -linear map on $\text{Hom}_{\mathcal{A}}(M, N)$ for all $M, N \in \text{mod } \mathcal{A}$ and $\delta(S(i)) \cong S(i)$ for all $i \in Q_0(\mathcal{A})$.


If there is an anti-automorphism ϵ of $\mathcal{A}$ (i.e., a K -map $\epsilon : \mathcal{A} \rightarrow \mathcal{A}$ with $\epsilon(a \cdot a') = \epsilon(a') \cdot \epsilon(a)$ and $\epsilon^2(a) = a$ for all $a, a' \in \mathcal{A}$) such that $\mathcal{A} \cdot \epsilon(e_i) \cong \mathcal{A} \cdot e_i$ for all $i \in Q_0(\mathcal{A})$, then $\mathcal{A}$ is a BGG-algebra (see [14]).

According to [9, Theorem 2.7], the quiver $Q(A)$ of a 1-quasi-hereditary algebra $(A, \leq)$ is the double of the quiver of the incidence algebra of $(\Lambda, \leq)$: Let $i, j \in \Lambda$, we write

$$i \triangleleft j \quad \text{and} \quad i \triangleright j$$

if i is a small neighbour of j and i is a large neighbour of j w.r.t. $\leq$, respectively. We have $\left| \left\{ \alpha \in Q_1(A) \mid i \xrightarrow{\alpha} j \right\} \right| = \begin{cases} 1 & \text{if } i \triangleleft j, \\ 1 & \text{if } i \triangleright j, \\ 0 & \text{else,} \end{cases}$ thus for any path $p =$

$(i_1 \rightarrow i_2 \rightarrow \dots \rightarrow i_m)$ there exists a uniquely determined path $\mathfrak{o}(p) := (i_m \rightarrow \dots \rightarrow i_2 \rightarrow i_1)$ in $Q(A)$ running through the same vertices in the opposite direction. Obviously, $\mathfrak{o}(\mathfrak{o}(p)) = p$.



1.3.1 Definition. A 1-quasi-hereditary algebra $A = KQ/\mathcal{I}$ is a $\text{BGG}_{(\trianglelefteq)}$ -algebra, if the K -linear map $\epsilon : KQ \rightarrow KQ$ given by $\epsilon(p) = \mathfrak{o}(p)$ for all paths p in Q induces an anti-automorphism of A .

Theorem B. Let $A \cong \text{End}_B(L)^{op} \cong KQ/\mathcal{I}$ with $(\Lambda, \leq)$ be a 1-quasi-hereditary algebra such that (B, L) satisfies the condition $\boxed{\leq}$. The following statements are equivalent:

(i) B is commutative.

(ii) A is a $\text{BGG}_{(\supseteq)}$ -algebra.

(iii) $\sum_{t=1}^r c_t \cdot p_t \in \mathcal{I}$ if and only if $\sum_{t=1}^r c_t \cdot \mathfrak{o}(p_t) \in \mathcal{I}$.


The Theorems A and B are special cases of Morita-Tachikawa Theorem 1.1. For the subsets $\mathbf{X}(1) := \{[A] \mid A \text{ is a 1-quasi-hereditary algebra}\}$, $\mathbf{X}' := \{[A] \mid A \text{ is a } \text{BGG}_{(\supseteq)}\text{-algebra}\}$

and

$\mathbf{Y}(1) := \{[B, L] \mid (B, L) \text{ has the property } \boxed{\leq}\}$, $\mathbf{Y}' := \{[B, L] \in \mathbf{Y}(1) \mid B \text{ is commutative}\}$

of $\mathbf{X}$ and $\mathbf{Y}$ respectively (defined in 1.1) we have $\Phi(\mathbf{X}(1)) = \mathbf{Y}(1)$ and $\Psi(\mathbf{Y}(1)) = \mathbf{X}(1)$ as well as $\Phi(\mathbf{X}') = \mathbf{Y}'$ and $\Psi(\mathbf{Y}') = \mathbf{X}'$. The function Φ restricted to $\mathbf{X}(1)$ maps $[A, \leq]$ to $[\text{End}_A(P(1))^{op}, P(1)]$, where $\{1\} = \min(\Lambda, \leq)$.

The picture to the right visualises this situation.




1.3.2 Example. Let $n \geq 3$ and $C = (c_{ij})_{2 \leq i, j \leq n-1} \in \text{GL}_{n-2}(K)$. We define $B := B_n(C) = K \langle x_2, \dots, x_{n-1} \rangle / \mathcal{I}$ with $\mathcal{I} := \langle \{c_{mk} \cdot x_i \cdot x_j - c_{ij} \cdot x_m \cdot x_k, x_i^3 \mid 2 \leq j, i, k, m \leq n-1\} \rangle$.

Let $X_k = x_k + \mathcal{I}$ for any $k \in \Gamma := \{2, \dots, n-1\}$. Since $\det C \neq 0$, for every $i \in \Gamma$ there exist $l(i), r(i) \in \Gamma$ such that $c_{l(i)i} \neq 0$ and $c_{ir(i)} \neq 0$. Therefore $c_{l(i)i} X_j X_{r(j)} = c_{jr(j)} X_{l(i)} X_i$ implies $\langle X_j X_{r(j)} \rangle = \langle X_{l(i)} X_i \rangle$ for all $i, j \in \Gamma$. Moreover, for any two $l_1(i), l_2(i) \in \Gamma$ with $c_{l_1(i)i} \neq 0$ and $c_{l_2(i)i} \neq 0$ we have $c_{l_1(i)i} X_{l_2(i)} X_i = c_{l_2(i)i} X_{l_1(i)} X_i$, hence $\langle X_{l_2(i)} X_i \rangle = \langle X_{l_1(i)} X_i \rangle$. A similar situation holds for any two $r_1(i), r_2(i) \in \Gamma$ with $c_{ir_1(i)} \neq 0$ and $c_{ir_2(i)} \neq 0$. Since $X_m^3 = 0$ for all $m \in \Gamma$, we obtain $X_i X_j X_k = 0$ for all $i, j, k \in \Gamma$.

Furthermore, we have $c_{ij} = 0$ iff $X_i X_j = 0$. Thus, $\langle X_i \rangle = B \cdot X_i = \text{span}_K \{X_i, X_{l(i)} X_i\} = \text{span}_K \{X_i, X_i X_{r(i)}\} = X_i \cdot B$ is a two-sided local ideal of B and $\text{soc } B = \langle X_j X_{r(j)} \rangle$ for all $j \in \Gamma$. The algebra B is self-injective and $\dim_K B = n$. Let $(\Lambda = \{1, \dots, n\}, \leq)$ be the poset given by $1 \triangleleft i \triangleleft n$ for all $2 \leq i \leq n-1$ and let the B -module $L := \bigoplus_{i=1}^n L(i)$ be given by $L(1) = B$, $L(i) = \langle X_i \rangle$ for all $i \in \Gamma$ and $L(n) = \text{soc } B$. The pair (B, L)

satisfies the property $\boxed{\leq}$. The quiver and relations of the 1-quasi-hereditary algebra $A_n(C) := \text{End}_B(L)^{op}$ can be found in [10, Example 3]. The algebra B is commutative if and only if $C = C^t$ and only in this case $A_n(C)$ is a $\text{BGG}_{(\supseteq)}$ -algebra.



2. Proof of the Theorem A

In this section let $(\Lambda, \leq)$ be a poset. Until the end of this paper for any $j \in \Lambda$ we denote by $\Lambda_{(j)}$ and $\Lambda^{(j)}$ the following subsets of Λ :

$$\Lambda_{(j)} := \{i \in \Lambda \mid i \leq j\} \quad \text{and} \quad \Lambda^{(j)} := \{i \in \Lambda \mid i \geq j\}.$$

We also adopt all notation of the previous section.

2.1. Proof of the Theorem A (i) $\Rightarrow$ (ii)

In this subsection A with $(\Lambda, \leq)$ denotes a 1-quasi-hereditary algebra and $\{1\} = \min(\Lambda, \leq)$. Furthermore, $B := \text{End}_A(P(1))^{op}$ and $L(i) := \text{Hom}_A(P(i), P(1))$ for any $i \in \Lambda$. Since $P(1)$ is a minimal faithful A -module, Theorem 1.1 and Lemma 1.2.3 provide isomorphisms

$$A \cong \text{End}_B({}_B P(1)) \quad \text{and} \quad {}_B P(1) \cong \bigoplus_{i \in \Lambda} L(i).$$

In particular, for any $[\mathcal{B}, \mathcal{L}] \in \mathbf{Y}$ with $A \cong \text{End}_{\mathcal{B}}(\mathcal{L})^{op}$ we obtain $\mathcal{B} \cong B$ and $\mathcal{L} \cong {}_B P(1)$. For the implication (i) $\Rightarrow$ (ii) we have to show that $(B, \bigoplus_{i \in \Lambda} L(i))$ satisfies the property $\boxed{\leq}$. We recall some notations and properties of 1-quasi-hereditary algebras from [9] and [10] which we will use in the proof: Let Q be the quiver of A and $\mathcal{I}$ be the corresponding ideal of KQ generated by the relations of A . The structure of Q (see 1.3) shows that for all $j, i, k \in \Lambda$ with $i \in \Lambda^{(j)} \cap \Lambda^{(k)}$ there exists a path $(j \rightarrow \lambda_1 \rightarrow \cdots \rightarrow \lambda_m \rightarrow i)$ with $j \leq \lambda_1 \leq \cdots \leq \lambda_m \leq i$ and a path $(i \rightarrow \mu_1 \rightarrow \cdots \rightarrow \mu_r \rightarrow k)$ with $i \geq \mu_1 \geq \cdots \geq \mu_r \geq k$. We write $p(j, i, i)$ resp. $p(i, i, k)$ for the residue class A of such a path. The concatenation of these two paths is denoted by $p(j, i, k) = p(i, i, k) \cdot p(j, i, i)$.

In the next subsection (Lemma 2.2.6) we will show, that for any two paths p and q in Q which yield paths of the form $p(j, i, k)$ in A we have $p - q \in \mathcal{I}$, thus we can talk about the path $p(j, i, k)$ in A . The A -map from $P(k)$ to $P(j)$ corresponding to $p(j, i, k)$ (i.e., $e_k \mapsto p(j, i, k)$) is denoted by $\mathfrak{f}_{(j,i,k)}$. For all j, i, k with $i \in \Lambda^{(j)} \cap \Lambda^{(k)}$ we have

$$\mathfrak{f}_{(j,i,k)} = \mathfrak{f}_{(j,i,i)} \circ \mathfrak{f}_{(i,i,k)} : \left(P(k) \xrightarrow{\mathfrak{f}_{(i,i,k)}} P(i) \xrightarrow{\mathfrak{f}_{(j,i,i)}} P(j) \right); \quad e_k \mapsto p(j, i, k) = p(i, i, k) \cdot p(j, i, i).$$

For any $i \in \Lambda^{(j)}$ the map $\mathfrak{f}_{(j,i,i)}$ is an inclusion and $\text{im}(\iota) = \text{im}(\mathfrak{f}_{(j,i,i)})$ for any inclusion $P(i) \xhookrightarrow{\iota} P(j)$ (see [9, 3.1(a)]). The contravariant functor $\text{Hom}_A(-, P(1)) : \text{mod } A \rightarrow \text{mod } B$ is exact since $P(1) \cong I(1)$ (see [9, 2.1(3)]). The inclusion $\mathfrak{f}_{(j,i,i)} : P(i) \hookrightarrow P(j)$ induces the surjective B -map $\underbrace{\text{Hom}_A(P(j), P(1))}_{L(j)} \twoheadrightarrow \underbrace{\text{Hom}_A(P(i), P(1))}_{L(i)}$ with $g \mapsto g \circ \mathfrak{f}_{(j,i,i)}$. Since

$P(i) \hookrightarrow P(j)$ if and only if $i \in \Lambda^{(j)}$ (see [9, 2.2]), we obtain $L(j) \twoheadrightarrow L(i)$ if and only if $j \leq i$, thus part (2)(a) of the Definition $\boxed{\leq}$ for $(B, \bigoplus_{i \in \Lambda} L(i))$ is satisfied. In particular, the surjection ${}_B B \cong L(1) \twoheadrightarrow L(i)$ with $F \mapsto F \circ \mathfrak{f}_{(1,i,i)}$ provides $L(i) = B \circ \mathfrak{f}_{(1,i,i)}$ for any $i \in \Lambda$.

It is enough to show the following two statements:

- ❶ The algebra B is local, self-injective and $\dim_K B = |\Lambda|$,
- ❷ $L(i) \cong B \circ \mathfrak{f}_{(1,i,1)} = \mathfrak{f}_{(1,i,1)} \circ B$ and $\text{rad}(B \circ \mathfrak{f}_{(1,i,1)}) = \sum_{i < i'} (B \circ \mathfrak{f}_{(1,i',1)})$ for any $i \in \Lambda$.

The second statement implies that $L(i)$ is isomorphic to a two-sided ideal of B for any $i \in \Lambda$ and $L(1) = {}_B B$ because $\mathfrak{f}_{(1,1,1)} = \text{id}_{P(1)}$. Moreover, we obtain the following explicit expression of the B -module $P(1)$:

2.1.1 Proposition. *For a 1-quasi-hereditary algebra $(A, \leq)$ with $\{1\} = \min(Q_0(A), \leq)$ and $B = \text{End}_A(P(1))^{op}$ we have ${}_B P(1) \cong \bigoplus_{i \in \Lambda} B \circ \mathfrak{f}_{(1,i,1)}$, where $\mathfrak{f}_{(1,i,1)}$ is the endomorphism of $P(1)$ as described above.*

2.1.2 Remark. The set $\{\mathbf{f}_{(j,i,k)} \mid i \in \Lambda^{(j)} \cap \Lambda^{(k)}\}$ is a K -basis of $\text{Hom}_A(P(k), P(j))$ for all $j, k \in \Lambda$ because the set $\{p(j, i, k) \mid i \in \Lambda^{(j)} \cap \Lambda^{(k)}\}$ is a K -basis of $P(j)_k$ (see [9, 3.2]). In particular, since $\Lambda^{(1)} = \Lambda$ we obtain that $\{\mathbf{f}_{(1,i,1)} \mid i \in \Lambda\}$ is a K -basis of B and

$\{\mathbf{f}_{(1,t,1)} \mid t \in \Lambda^{(i)}\}$ is a K -basis of $L(i) = \text{Hom}_A(P(i), P(1))$ for any $i \in \Lambda$.

This situation gives rise to some implications: Let $f \in \text{End}_A(P(j))^{op}$ and $i \in \Lambda^{(j)}$, then $f \circ \mathbf{f}_{(j,i,i)} \in \text{Hom}_A(P(i), P(j)) = \text{span}_K \{\mathbf{f}_{(j,t,i)} \mid t \in \Lambda^{(i)}\}$, since $\Lambda^{(j)} \cap \Lambda^{(i)} = \Lambda^{(i)}$.


Thus $f \circ \mathbf{f}_{(j,i,i)} = \sum_{t \in \Lambda^{(i)}} c_t \cdot \mathbf{f}_{(j,t,i)}$ for $c_t \in K$. Let $k \in \Lambda_{(i)}$, then

$$p(j, t, k) = p(i, i, k) \cdot p(j, t, i) \text{ implies } \mathbf{f}_{(j,t,k)} = \mathbf{f}_{(j,t,i)} \circ \mathbf{f}_{(i,i,k)}.$$

We obtain $f \circ \mathbf{f}_{(j,i,k)} = (f \circ \mathbf{f}_{(j,i,i)}) \circ \mathbf{f}_{(i,i,k)} = \sum_{t \in \Lambda^{(i)}} c_t \cdot \mathbf{f}_{(j,t,k)}$ for every $t \in \Lambda^{(i)}$.

In other words, for any $i \in \Lambda^{(j)} \cap \Lambda^{(k)}$ and $f \in \text{End}_A(P(j))^{op}$ we have

$f \circ \mathbf{f}_{(j,i,k)} \in \text{span}_K \{\mathbf{f}_{(j,t,k)} \mid t \in \Lambda^{(i)}\}$. In particular, $B \circ \mathbf{f}_{(1,i,k)} \subseteq \text{span}_K \{\mathbf{f}_{(1,t,k)} \mid t \in \Lambda^{(i)}\}$.



Proof of ❶. The algebra A^{op} is also 1-quasi-hereditary with $P_{A^{op}}(1)$ as a minimal faithful A^{op} -module (see [9, 1.3]). The algebras $B^{op} \cong \text{End}_A(P(1))$ and $B \cong \text{End}_{A^{op}}(P_{A^{op}}(1)) \cong \text{End}_A(P(1))^{op}$ are local, since $P(1)$ and $P_{A^{op}}(1)$ are indecomposable. Therefore $\text{soc } B$ and $\text{top } B$ are simple, thus B is local, self-injective and $\dim_K B = |\{\mathbf{f}_{(1,i,1)} \mid i \in \Lambda\}| = |\Lambda|$. $\square$


The proof of ❷ is based on the following properties of B -modules generated by the maps $\mathbf{f}_{(1,i,k)}$.

2.1.3 Lemma. For all $i, k \in \Lambda$ with $i \in \Lambda^{(k)}$ the following is satisfied:

- (1) $B \circ \mathbf{f}_{(1,i,k)} = \text{span}_K \{\mathbf{f}_{(1,t,k)} \mid t \in \Lambda^{(i)}\}$ for all $k \in \Lambda_{(i)}$,
- (2) $B \circ \mathbf{f}_{(1,i,k)} \cong B \circ \mathbf{f}_{(1,i,k')}$ for all $k, k' \in \Lambda_{(i)}$,
- (3) $B \circ \mathbf{f}_{(1,i,1)} = \mathbf{f}_{(1,i,1)} \circ B = \text{span}_K \{\mathbf{f}_{(1,t,1)} \mid t \in \Lambda^{(i)}\}$.

We will clarify the statements of this lemma using an example: Let A be the algebra corresponding to a regular block of $\mathcal{O}(\mathfrak{sl}_3(\mathbb{C}))$, then $P(1)$ is the minimal faithful A -module (the presentation of $P(1)$ via the $\mathbb{C}$ -basis $p(1, i, k)$ can be found in [10, Sec.2]).

The picture presents $P(1)$ as a $\text{End}_A(P(1))^{op}$ -module (in particular, $\text{End}_A(P(1))^{op} \cong \mathbb{C}[x, y] / \langle xy, x^3 - y^3 \rangle$, see Example 1.2): The circle $C(k)$ represents the spaces $L(k) = \text{Hom}_A(P(j), P(1))$ for every $k \in \Lambda = \{1, \dots, 6\}$. Inside of $C(k)$ we find the $\mathbb{C}$ -basis $\{\mathbf{f}_{(i,k)} := \mathbf{f}_{(1,i,k)} \mid i \in \Lambda^{(k)}\}$ of $L(k)$. There is an arrow $\mathbf{f}_{(t,k)} \rightarrow \mathbf{f}_{(i,k)}$ if $B \circ \mathbf{f}_{(t,k)} \subset B \circ \mathbf{f}_{(i,k)}$ (statement (1)) and an arrow $\mathbf{f}_{(i,k)} \dashrightarrow \mathbf{f}_{(i,k')}$ if $B \circ \mathbf{f}_{(i,k)} \cong B \circ \mathbf{f}_{(i,k')}$ (statement (2)). We focus on the B -submodules of $L(k)$ generated by $\mathbf{f}_{(4,k)}$ for $k \in \Lambda_{(4)} = \{1, 2, 3, 4\}$. Statement (1) yields $B \circ \mathbf{f}_{(4,k)} = \text{span}_K \{\mathbf{f}_{(4,k)}, \mathbf{f}_{(6,k)}\}$ (visualized in the parallelogram). Statement (2) implies $L(4) \cong B \circ \mathbf{f}_{(4,k)}$ for all $k \in \Lambda_{(4)}$. The B -module $L(4)$ is isomorphic to the two-sided ideal $B \circ \mathbf{f}_{(4,1)} = \mathbf{f}_{(4,1)} \circ B$ of B since $\mathbf{f}_{(4,1)} \in B$ (statement (3)).



Proof. (1) Since $P(1) \cong I(1)$ and $\mathfrak{f}_{(1,i,i)} : P(i) \hookrightarrow P(1)$, the universal property of injective modules imply that for any $t \in \Lambda^{(i)}$ there exists $F(t) \in B$ such that $F(t) \circ \mathfrak{f}_{(1,i,i)} = \mathfrak{f}_{(1,t,i)}$. Let $k \in \Lambda(i)$, then $\mathfrak{f}_{(1,i,i)} \circ \mathfrak{f}_{(i,i,k)} = \mathfrak{f}_{(1,i,k)}$ provides the commutative diagram

$$\begin{array}{ccccc} \mathfrak{f}_{(1,t,k)} : & P(k) & \xrightarrow{\mathfrak{f}_{(i,i,k)}} & P(i) & \xrightarrow{\mathfrak{f}_{(1,t,i)}} & P(1) \\ & \parallel & & \downarrow \mathfrak{f}_{(1,i,i)} & & \parallel \\ F(t) \circ \mathfrak{f}_{(1,i,k)} : & P(k) & \xrightarrow{\mathfrak{f}_{(1,i,k)}} & P(1) & \xrightarrow{F(t)} & P(1) \end{array}$$

We obtain $F(t) \circ \mathfrak{f}_{(1,i,k)} = \mathfrak{f}_{(1,t,i)} \circ \mathfrak{f}_{(i,i,k)} = \mathfrak{f}_{(1,t,k)}$. In other words, we have $B \circ \mathfrak{f}_{(1,i,k)} \supseteq \text{span}_K \{ \mathfrak{f}_{(1,t,k)} \mid t \in \Lambda^{(i)} \}$. The previous remark yields $B \circ \mathfrak{f}_{(1,i,k)} \subseteq \text{span}_K \{ \mathfrak{f}_{(1,t,k)} \mid t \in \Lambda^{(i)} \}$.

(2) Let $k \in \Lambda(i)$. We consider the map $(-\circ \mathfrak{f}_{(k,k,1)}) : B \circ \mathfrak{f}_{(1,i,k)} \rightarrow B \circ \mathfrak{f}_{(1,i,1)}$, and we have $F \circ \mathfrak{f}_{(1,i,k)} \mapsto F \circ \mathfrak{f}_{(1,i,k)} \circ \mathfrak{f}_{(k,k,1)} = F \circ \mathfrak{f}_{(1,i,1)}$ (see 2.1.2). Obviously, this map is a surjective B -map. Moreover, $\dim_K (B \circ \mathfrak{f}_{(1,i,1)}) \stackrel{(1)}{=} \dim_K (B \circ \mathfrak{f}_{(1,i,k)})$ since $\{ \mathfrak{f}_{(1,t,k)} \mid t \in \Lambda^{(i)} \}$ is K -independent. Thus $B \circ \mathfrak{f}_{(1,i,1)} \cong B \circ \mathfrak{f}_{(1,i,k)}$ for all $k \in \Lambda(i)$.

(3) It is enough to show $\mathfrak{f}_{(1,i,1)} \circ B = \text{span}_K \{ \mathfrak{f}_{(1,t,1)} \mid t \in \Lambda^{(i)} \}$ [the rest follows from (1)]: For any $i \in \Lambda$ we have $\mathfrak{f}_{(1,i,1)} = \mathfrak{f}_{(1,i,i)} \circ \mathfrak{f}_{(i,i,1)}$ and $\mathfrak{f}_{(i,i,1)} \circ F \in \text{Hom}_A(P(1), P(i)) = \text{span}_K \{ \mathfrak{f}_{(i,t,1)} \mid t \in \Lambda^{(i)} \}$ for all $F \in B$. With similar arguments as in the previous remark we obtain $\mathfrak{f}_{(1,t,1)} = \mathfrak{f}_{(1,i,i)} \circ \mathfrak{f}_{(i,t,1)}$. This provides $\mathfrak{f}_{(1,i,1)} \circ F \in \text{span}_K \{ \mathfrak{f}_{(1,t,1)} \mid t \in \Lambda^{(i)} \}$ for every $F \in B$. Therefore, we obtain $\mathfrak{f}_{(1,i,1)} \circ B \subseteq \text{span}_K \{ \mathfrak{f}_{(1,t,1)} \mid t \in \Lambda^{(i)} \}$.

For any $t \in \Lambda^{(i)}$ the A -module generated by $p(1, t, 1)$ is a submodule of $A \cdot p(1, i, 1)$ (see [10, 2.2 (a)]). There exists $p(t) \in P(1)_1$ with $p(1, t, 1) = p(t) \cdot p(1, i, 1)$. Let $F(t) \in B$ be given by $F(t)(e_1) = p(t)$, then $\mathfrak{f}_{(1,t,1)} = \mathfrak{f}_{(1,i,1)} \circ F(t)$. Thus $\mathfrak{f}_{(1,i,1)} \circ B \supseteq \text{span}_K \{ \mathfrak{f}_{(1,t,1)} \mid t \in \Lambda^{(i)} \}$. $\square$

Proof of 2. The parts (2) and (3) of the previous lemma imply $L(i) = B \circ \mathfrak{f}_{(1,i,i)} \cong B \circ \mathfrak{f}_{(1,i,1)} = \mathfrak{f}_{(1,i,1)} \circ B$. Thus $L(i)$ is a two-sided local ideal of B for all $i \in \Lambda$.

The B -module $B \circ \mathfrak{f}_{(1,i,1)}$ is local, thus $B \circ \mathfrak{f}_{(1,t,1)} \stackrel{(1)}{\subset} B \circ \mathfrak{f}_{(1,i,1)}$ for all $t \in \Lambda^{(i)} \setminus \{i\} = \{t \in \Lambda \mid i < t\}$ implies $\sum_{i < t} (B \circ \mathfrak{f}_{(1,t,1)}) \subseteq \text{rad}(B \circ \mathfrak{f}_{(1,i,1)})$. Since $\{ \mathfrak{f}_{(1,t,1)} \mid i < t \}$ is linearly independent and $\{ \mathfrak{f}_{(1,t,1)} \mid i < t \} \subseteq \sum_{i < t} (B \circ \mathfrak{f}_{(1,t,1)})$, we have $\dim_K (\sum_{i < t} (B \circ \mathfrak{f}_{(1,t,1)})) \geq |\Lambda^{(i)}| - 1 \stackrel{(1)}{=} \dim_K (B \circ \mathfrak{f}_{(1,i,1)}) - 1 = \dim_K \text{rad}(B \circ \mathfrak{f}_{(1,i,1)})$. We obtain $\sum_{i < t} (B \circ \mathfrak{f}_{(1,t,1)}) = \text{rad}(B \circ \mathfrak{f}_{(1,i,1)})$ for all $i \in \Lambda$. $\square$

2.2. Proof of the Theorem A (ii) $\Rightarrow$ (i)

Let B be a local, self-injective K -algebra with $\dim_K B = n$. Let the set $(\Lambda = \{1, \dots, n\}, \leq)$ be partially ordered and let $L \cong \bigoplus_{i \in \Lambda} L(i)$ be a B -module, such that (B, L) satisfies the condition $\boxed{\leq}$ (see Definition 1.2.4).

It is easy to see that there exists a uniquely determined minimal and maximal element in $(\Lambda, \leq)$: Since $B (= L(1))$ is a projective cover of any local B -module, we have $L(1) \twoheadrightarrow L(i)$, thus $\boxed{\leq}$ (2)(a) implies $1 \leq i$ for all $i \in \Lambda$. Let $k \in \Lambda$ be maximal, then $\boxed{\leq}$ (2) (b) yields $\text{rad } L(k) = 0$. The local submodule $L(k)$ of B is simple, thus $L(k) = \text{soc}(B)$, because the socle of a local self-injective algebra is simple. Since $L(j) \cong L(i)$ iff $j = i$ (see $\boxed{\leq}$ (2)(a)), we have $L(k) \subseteq \text{rad } L(i) \stackrel{(b)}{=} \sum_{i < j} L(j)$ for every $i \in \Lambda$ with $i \neq k$. Consequently, $i \leq k$ for all

$i \in \Lambda$. The maximal element in $(\Lambda, \leq)$ will be denoted by $n = |\Lambda|$. We have $\{1\} = \min(\Lambda, \leq)$ and $\{n\} = \max(\Lambda, \leq)$ with $L(1) = B$ and $L(n) = \text{soc}(B)$.

We consider the algebra $A := \text{End}_B(L)^{\text{op}}$. Because L is multiplicity-free and has n direct summands, A is basic and the quiver $Q(A)$ has n vertices. We identify these with the elements in Λ . In order to prove that $(A, \leq)$ is a 1-quasi-hereditary algebra (see Definition 1.2.2) we have to show that for all $j \in \Lambda$ the following holds:

- ❶ $[\Delta(j) : S(i)] = 1$ for all $i \in \Lambda_{(j)}$,
- ❷ $P(j)$ has a Δ -good filtration with $(P(j) : \Delta(i)) = \begin{cases} 1 & \text{if } i \in \Lambda^{(j)}, \\ 0 & \text{else,} \end{cases}$
- ❸ $\text{soc } P(j) \cong \text{top } I(j) \cong S(1)$,
- ❹ $\Delta(j) \hookrightarrow \Delta(n)$ and $\nabla(n) \twoheadrightarrow \nabla(j)$.

Recall that an (left) ideal J of a local, self-injective (basic) algebra $\mathcal{B}$ is local if and only if it is generated by some non-zero element $x \in \mathcal{B}$ (i.e., $J = \mathcal{B} \cdot x$). Moreover, $\dim_K J = \dim_K(\text{rad } J) + 1$ and $J = \text{span}_K \{x \cup \text{rad } J\}$. In addition, for an ideal M of $\mathcal{B}$, any $\mathcal{B}$ -map $f : J \rightarrow M$ is induced by right-multiplication by an element $b_f \in \mathcal{B}$, more precisely, $f(y \cdot x) = y \cdot x \cdot b_f$ for all $y \in \mathcal{B}$ (in this case we write $J \xrightarrow{b_f} M$). The element $f(x) = x \cdot b_f$ generates $\text{im}(f) = \mathcal{B} \cdot f(x)$. In particular, we have $\text{Hom}_{\mathcal{B}}(J, \mathcal{B}) = \{J \xrightarrow{b} \mathcal{B} \mid b \in \mathcal{B}\}$.

The annihilator of $L(j)$ is $\text{Ann}(L(j)) := \{b \in B \mid b \cdot L(j) = 0\}$.

2.2.1 Lemma. *Let B be a local, self-injective algebra and let $(\Lambda, \leq)$ be a poset with $\dim_K B = |\Lambda| = n$. Let $L \cong \bigoplus_{i \in \Lambda} L(i)$ be a B -module such that (B, L) satisfies the condition*

$\boxed{\leq}$. *Then for all $i, j, k \in \Lambda$ the following properties are satisfied:*

(1) *Let $x_i \in B$ be a generator of $L(i)$ and $X(i) := \{x_k \mid k \in \Lambda^{(i)}\}$. Then we have:*

(1.1) *The set $X(i)$ is a K -basis of $L(i)$. In particular, for any subset $\Gamma \subseteq \Lambda$ the set $\bigcup_{j \in \Gamma} X(j)$ is a K -basis of $\sum_{j \in \Gamma} L(j)$ and $\bigcap_{j \in \Gamma} X(j)$ is a K -basis of $\bigcap_{j \in \Gamma} L(j)$.*

(1.2) $L(i) = B \cdot x_i = x_i \cdot B$.

(2) *Let N be a submodule of B , then $\text{im}(f) \subseteq L(i) \cap N$ for all $f \in \text{Hom}_B(L(i), N)$.*

(3) *We have $L(i) \hookrightarrow L(j)$ resp. $L(j) \twoheadrightarrow L(i)$ if and only if $i \in \Lambda^{(j)}$. Moreover,*

(3.1) $\text{im}(L(i) \hookrightarrow L(j)) = L(i)$ for any injective B -map from $L(i)$ to $L(j)$,

(3.2) $\ker(L(j) \twoheadrightarrow L(i)) = \text{Ann}(L(i)) \cdot L(j)$ for any surjective B -map from $L(j)$ to $L(i)$.

(4) *For $i \in \Lambda^{(j)} \cap \Lambda^{(k)}$ let $f_{(k,i,i)} : L(i) \hookrightarrow L(k)$ be an injective, $f_{(i,i,j)} : L(j) \twoheadrightarrow L(i)$ a surjective B -map and $f_{(k,i,j)} := f_{(k,i,i)} \circ f_{(i,i,j)} (\in \text{Hom}_B(L(j), L(k)))$. The set*

$$\mathfrak{B}(k, j) := \{f_{(k,i,j)} \mid i \in \Lambda^{(j)} \cap \Lambda^{(k)}\} \text{ is a } K\text{-basis of } \text{Hom}_B(L(j), L(k)).$$

In particular, every map $f \in \text{Hom}_B(L(j), L(k))$ factors through $\bigoplus_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} L(i)$.

Proof. (1.1) By induction on $\dim_K(L(j))$ we show that the K -space $L(j)$ is spanned by $X(j)$ for every $j \in \Lambda$: If $\dim_K(L(j)) = 1$, then $L(j)$ is simple, thus $j = n$ and $X(n) = \{x_n\}$ is a K -basis of $L(n) = \text{soc } B$. Let $j \in \Lambda$ with $\dim_K(L(j)) = m + 1$, then $L(l) = \text{span}_K X(l)$ for any $l \in \Lambda^{(j)} \setminus \{j\}$, because $L(l) \subseteq \text{rad } L(j)$ (see 1.2.4(2)(b)) implies $\dim_K L(l) \leq \dim_K(\text{rad } L(j)) = m$. Thus $\text{rad } L(j) = \sum_{j < l} L(l)$ is spanned by $\bigcup_{j < l} X(l) = \{x_k \mid j < k\}$ and consequently $X(j) = \{x_j\} \cup \bigcup_{j < l} X(l)$ spans the K -space $L(j)$. Furthermore, $B = L(1) = \text{span}_K X(1) = \{x_1, \dots, x_n\}$ and $\dim_K B = n$ implies that $X(1)$ is a K -basis of B and consequently the subset $X(j)$ of $X(1)$ is linearly independent.

For $\Gamma \subseteq \Lambda$ the subsets $\bigcup_{j \in \Gamma} X(j) = \{x_k \mid k \in \Lambda, k \geq i \text{ for some } i \in \Gamma\}$ and $\bigcap_{j \in \Gamma} X(j) = \{x_k \mid k \in \bigcap_{j \in \Gamma} \Lambda^{(j)}\}$ of $X(1)$ generate $\sum_{j \in \Gamma} L(j)$ and $\bigcap_{j \in \Gamma} L(j)$ as K -spaces respectively.

(1.2) Since $L(j) \twoheadrightarrow L(i)$ for any $i \in \Lambda^{(j)}$ (see 1.2.4(2)(a)), there exists $b_i \in B$ with $L(j) \xrightarrow{b_i} L(i)$ and $x_i = x_j \cdot b_i$. Let $y \in L(j) = B \cdot x_j$, then $y \stackrel{(1.1)}{=} \sum_{i \in \Lambda^{(j)}} c_i \cdot x_i = \sum_{i \in \Lambda^{(j)}} c_i \cdot x_j \cdot b_i = x_j \cdot (\sum_{i \in \Lambda^{(j)}} c_i \cdot b_i)$ (here $c_i \in K$). We obtain $B \cdot x_j \subseteq x_j \cdot B$. Since $L(j)$ is a two-sided ideal, we have $B \cdot x_j \cdot B \subseteq B \cdot x_j$, thus $x_j \cdot B \subseteq B \cdot x_j$ and consequently $L(j) = B \cdot x_j = x_j \cdot B$.

(2) For $f \in \text{Hom}_B(L(i), N)$ there exists some $b \in B$ with $f : L(i) \xrightarrow{b} N$. Let x_i be a generator of $L(i)$, then $f(x_i) = x_i \cdot b = \tilde{b} \cdot x_i$ for some $\tilde{b} \in B$ by (1.2). Thus $f(x_i) \in L(i)$ and consequently $\text{im}(f) \subseteq L(i) \cap N$.

(3.1) According to 1.2.4(2)(b) we have $L(i) \subseteq L(k)$ if and only if $i \in \Lambda^{(k)}$. Let N be a submodule of $L(k)$ with $N \cong L(i)$ and $f \in \text{Hom}_B(L(i), N)$ be an isomorphism, then $f(L(i)) = N \subseteq L(i) \cap N$ (see (2)). We obtain $f(L(i)) = L(i) = N$.

(3.2) Let $\pi_i : L(j) \twoheadrightarrow L(i)$ be a surjection and x_j a generator of $L(j)$. Then $x_i := \pi_i(x_j)$ generates $L(i)$. Let $x \in L(j)$, then $x = b \cdot x_j$ for some $b \in B$. Obviously, $x \in \ker(\pi_i)$ if and only if $\pi_i(x) = b \cdot x_i = 0$ if and only if $b \cdot x_i \cdot B \stackrel{(1.2)}{=} b \cdot L(i) = 0$. We obtain $\ker(\pi_i) = \{b \cdot x_j \in L(j) \mid b \in \text{Ann}(L(i))\} = \text{Ann}(L(i)) \cdot L(j)$.

(4) Let x_j be some generator of $L(j)$. Then $x_i := f_{(k,i,j)}(x_j)$ generates the submodule $L(i)$ of $L(j) \cap L(k)$. The set $\{x_i \mid i \in \Lambda^{(j)} \cap \Lambda^{(k)}\} \stackrel{(1.1)}{=} X(j) \cap X(k)$ is a K -basis of $L(j) \cap L(k)$. Let $f \in \text{Hom}_B(L(j), L(k))$, then $\text{im}(f) \subseteq L(j) \cap L(k)$ (see (2)). Thus $f(x_j) = \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} c_i \cdot x_i$ and consequently $f = \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} c_i \cdot f_{(k,i,j)}$. Let f_1, f_2 be the B -maps given by $f_1 : L(j) \rightarrow \bigoplus_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} L(i)$ with $f_1(x_j) = (\pi_i(x_j))_{i=1, \dots, n}$ and $f_2 : \bigoplus_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} L(i) \rightarrow L(k)$ with $f_2(x_i)_{i=1, \dots, n} = \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} c_i \cdot x_i$. We have $f = f_2 \circ f_1$ and thus f factors through $\bigoplus_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} L(i)$. $\square$

2.2.2 Remark. For a pair (B, L) satisfying the property $\boxed{\leq}$ with $L = \bigoplus_{i \in \Lambda} L(i)$ denote by B the algebra B^{op} and by $L(i)$ the B -module $\mathcal{D}(L(i))$, where $\mathcal{D} : \text{mod } B \rightarrow \text{mod } B$ is the standard duality functor. Since $\text{top } L(i)$ and $\text{soc } L(i)$ are simple, we obtain that $\text{soc } L(i)$ and $\text{top } L(i)$ are simple for all $i \in \Lambda$. In particular, $B (= L(1))$ is a local, self-injective algebra with $\dim_K B = \dim_K B = n$ and $L(i)$ can be considered as a local (left) ideal of B for any $i \in \Lambda$.

We denote by $\mathbf{f}_{(j,i,k)}$ the B -map $\mathcal{D}(f_{(k,i,j)}) : L(k) \rightarrow L(j)$ for all $i, j, k \in \Lambda$ with $i \in \Lambda^{(j)} \cap \Lambda^{(k)}$, where $f_{(k,i,j)}$ is the B -map described in 2.2.1(4). As $\mathcal{D}$ is duality, we have $\mathbf{f}_{(j,i,k)} : L(k) \xrightarrow{\mathbf{f}_{(i,i,k)}} L(i) \xrightarrow{\mathbf{f}_{(j,i,i)}} L(j)$.

2.2.3 Proposition. A pair (B, L) satisfies the condition $\boxed{\leq}$ if and only if $(B^{op}, \mathcal{D}(L))$ satisfies the condition $\boxed{\leq}$.

Proof. (We use the notations introduced in 2.2.2.) According to 2.2.1 (3) we have $L(j) \twoheadrightarrow L(i)$ resp. $L(i) \hookrightarrow L(j)$ if and only if $i \in \Lambda^{(j)}$. Moreover, any two surjections $\tilde{\pi}_1, \tilde{\pi}_2$ from $L(j)$ to $L(i)$ are induced by some injective maps $\iota_1, \iota_2 : L(i) \hookrightarrow L(j)$ and $\ker(\tilde{\pi}_k) = \{\xi \in \text{Hom}_K(L(j), K) \mid \xi|_{\text{im}(\iota_k)} = 0\}$, here $k = 1, 2$. Since $\text{im}(\iota_1) \stackrel{2.2.1(3)}{=} \text{im}(\iota_2)$, we obtain $\ker(\tilde{\pi}_1) = \ker(\tilde{\pi}_2)$. Similarly, $\text{im}(\tilde{\iota}) = L(i)$ for all injections $\tilde{\iota} : L(i) \hookrightarrow L(j)$.

Let $1 := 1_B$ and $y_i := f_{(j,i,1)}(1)$ for any $i \in \Lambda^{(j)}$. Obviously, y_i is a generator of the submodule $L(i)$ of $L(j)$. According to 2.2.1(4) the set $\{f_{(j,i,1)} \mid i \in \Lambda^{(j)}\}$ is a K -basis of $\text{Hom}_B(L(1), L(j))$. Thus the set $X(j) := \{y_i \mid i \in \Lambda^{(j)}\}$ is a K -basis of $L(j)$, since $\dim_K L(j) = \dim_K L(j) \stackrel{2.2.1(1)}{=} |X(j)|$.

Now we show $L(j) = B \cdot y_j = y_j \cdot B$ (this implies, that $L(j)$ is a two-sided ideal of B):

Let $f_{(i,i,j)} : L(j) \xrightarrow{b_i} L(i)$ such that $y_i = y_j \cdot b_i$ for any $i \in \Lambda^{(j)}$. Let $y \in B \cdot y_j$, then $y = \sum_{i \in \Lambda^{(j)}} c_i \cdot y_i = y_j \cdot (\sum_{i \in \Lambda^{(j)}} c_i \cdot b_i) \in y_j \cdot B$, thus $B \cdot y_j \subseteq y_j \cdot B$. On the other hand, for some $y \in y_j \cdot B$ with $y = y_j \cdot b$ we have $y \in \text{im}(f)$, where $f \in \text{Hom}_B(L(j), L(1)) = \text{span}_K \{f_{(1,i,j)} \mid i \in \Lambda^{(j)}\}$ is given by $f : L(j) \xrightarrow{b} L(1) = B$. Since $\text{im}(f_{(1,i,j)}) = L(i) \subseteq L(j)$ for each $i \in \Lambda^{(j)}$, we obtain $\text{im}(f) \subseteq L(j)$ and consequently $y \in L(j) = B \cdot y_j$. Thus we obtain $B \cdot y_j \supseteq y_j \cdot B$.

If $i \neq j$, then $L(i) \neq L(j)$, thus $L(i) \subseteq \text{rad}(L(j))$ for any $i \in \Lambda^{(j)} \setminus \{j\}$. Consequently, $\sum_{j < i} L(i) \subseteq \text{rad}(L(j))$. The set $\bigcup_{j < i} X(i) = \{y_i \mid i \in \Lambda^{(j)} \setminus \{j\}\}$ is a K -basis of $\sum_{j < i} L(i)$, since $X(i)$ is a K -basis of $L(i)$. Thus $\dim_K \left(\sum_{j < i} L(i) \right) = |\Lambda^{(j)} \setminus \{j\}| = |\Lambda^{(j)}| - 1 = \dim_K L(j) - 1 = \dim_K (\text{rad } L(j))$ implies $\sum_{j < i} L(i) = \text{rad}(L(j))$ for all $j \in \Lambda$. $\square$

Furthermore, for a pair (B, L) which satisfies the condition $\boxed{\leq}$ we consider the algebra $A = \text{End}_B(L) \cong \text{End}_B(L)^{op}$. The evaluation functor $\text{Hom}_B(L, -) : \text{mod } B \rightarrow \text{mod } A$ provides an isomorphism $\text{Hom}_B(L(i), L(j)) \cong \text{Hom}_A(P(i), P(j))$ (see [1, Proposition 2.1]). Moreover, an injective B -map $L(i) \xhookrightarrow{f} L(j)$ induces an injective A -map $P(i) \xhookrightarrow{f} P(j)$, since $\text{Hom}_B(L, -)$ is left exact.

The previous Lemma shows that the properties described in Lemma 2.2.1 are also satisfied for the B -ideals $L(i)$. The part (3) implies $P(i) \hookrightarrow P(j)$ for all $i \in \Lambda^{(j)}$, moreover, since $\text{im}(L(i) \hookrightarrow L(j)) = L(i)$ for any injective B -map, we obtain that a submodule of $P(j)$ isomorphic to $P(i)$ is uniquely determined. For $i \in \Lambda^{(j)}$ we consider $P(i)$ as a submodule of $P(j)$.

For all $i \in \Lambda^{(j)} \cap \Lambda^{(k)}$ let $f_{(j,i,k)} \in \text{Hom}_A(P(k), P(j))$ be the map induced by $f_{(j,i,k)} \in \text{Hom}_B(L(k), L(j))$ (described in 2.2.2). Since $f_{(j,i,k)} = f_{(j,i,i)} \circ f_{(i,i,k)}$ and $f_{(j,i,i)}$ is injective, we obtain $f_{(j,i,k)} = f_{(j,i,i)} \circ f_{(i,i,k)} : \left(P(k) \xrightarrow{f_{(i,i,k)}} P(i) \xhookrightarrow{f_{(j,i,i)}} P(j) \right)$. Obviously, $\text{im}(f_{(j,i,k)})$ belongs to the submodule $P(i)$ of $P(j)$.

2.2.4 Remark-Notations. According to Lemma 2.2.1(4) the set $\{f_{(j,i,k)} \mid i \in \Lambda^{(j)} \cap \Lambda^{(k)}\}$ is a K -basis of $\text{Hom}_A(P(k), P(j))$ for all $j, k \in \Lambda$. Thus $\text{im}(f) \subseteq \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} \text{im}(f_{(j,i,k)}) \subseteq \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} P(i)$ for all $f \in \text{Hom}_A(P(k), P(j))$. Let $p(j, i, k) := f_{(j,i,k)}(e_j)$, then we obtain that $\{p(j, i, k) \mid i \in \Lambda^{(j)} \cap \Lambda^{(k)}\}$ is a K -basis of $P(j)_k$.

Let Γ be some subset of $\Lambda^{(j)}$. Then the following hold:

- (1) For any $l \in \Lambda$ the set $\mathbf{B}_j(\Gamma, l) := \{p(j, t, l) \mid t \in \bigcup_{i \in \Gamma} (\Lambda^{(i)} \cap \Lambda^{(l)})\}$ is a K -basis of the subspace $(\sum_{i \in \Gamma} P(i))_l$ of $P(j)_l$ for the submodule $\sum_{i \in \Gamma} P(i)$ of $P(j)$: Let $i \in \Gamma$, then $\{f_{(i, t, l)} \mid t \in \Lambda^{(i)} \cap \Lambda^{(l)}\}$ is a K -basis of $\text{Hom}_A(P(l), P(i))$. By applying $\text{Hom}_A(P(l), -)$ to $f_{(j, i, i)} : P(i) \hookrightarrow P(j)$, $e_i \mapsto p(j, i, i)$, we obtain $\text{Hom}_A(P(l), P(i)) \hookrightarrow \text{Hom}_A(P(l), P(j))$ with $f_{(i, t, l)} \mapsto f_{(j, i, i)} \circ f_{(i, t, l)} = f_{(j, t, l)}$ (or, equivalently, $P(i)_l \hookrightarrow P(j)_l$ with $p(i, t, l) \mapsto p(j, t, l)$). The set $\{p(j, t, l) \mid t \in \Lambda^{(i)} \cap \Lambda^{(l)}\}$ is a K -basis of $P(i)_l \subseteq P(j)_l$ for any $i \in \Gamma$. Thus $\mathbf{B}_j(\Gamma, l) = \bigcup_{i \in \Gamma} \{p(j, t, l) \mid t \in \Lambda^{(i)} \cap \Lambda^{(l)}\}$ is a K -basis of the subspace $(\sum_{i \in \Gamma} P(i))_l$ of $P(j)_l$.
- (2) The set $\mathbf{B}_j(\Gamma) := \{p(j, t, l) \mid t \in \bigcup_{i \in \Gamma} \Lambda^{(i)}, l \in \Lambda_{(t)}\}$ is a K -basis of the submodule $\sum_{i \in \Gamma} P(i)$ of $P(j)$: Obviously, $\mathbf{B}_j(\Gamma)$ is the disjoint union of $\mathbf{B}_j(\Gamma, l)$ for $l \in \Lambda$, thus $\mathbf{B}_j(\Gamma)$ is a K -basis of $\sum_{i \in \Gamma} P(i)$.

We can now prove the four statements formulated at the beginning of this subsection, so that the algebra A with $(\Lambda, \leq)$ is 1-quasi-hereditary (recall that $1 \leq i \leq n$ for all $i \in \Lambda$).

Proof. ❶ The definition of the standard modules provides $\Delta(j) = P(j)/N(j)$, where $N(j) = \sum_{k \not\leq j} \sum_{f \in \text{Hom}_A(P(k), P(j))} \text{im}(f)$. The previous deliberations imply $N(j) \subseteq \sum_{j < i} P(i)$. Since $P(i) = \text{im}(f_{(j, i, i)}) \subseteq N(j)$ for any $i \in \Lambda^{(j)} \setminus \{j\}$, we obtain $N(j) \supseteq \sum_{j < i} P(i)$, thus $\Delta(j) = P(j) / \left(\sum_{j < i} P(i) \right)$.

Let $\Gamma = \Lambda^{(j)} \setminus \{j\}$ and $k \in \Lambda_{(j)}$, then $\mathbf{B}_j(\Lambda^{(j)}, k) = \{p(j, j, k)\} \dot{\cup} \mathbf{B}_j(\Gamma, k)$, using the notation of 2.2.4(1). Since $P(j)_k = \text{span}_K \mathbf{B}(\Lambda^{(j)}, k)$ and $\left(\sum_{j < i} P(i) \right)_k = \text{span}_K \mathbf{B}_j(\Gamma, k)$, we obtain $1 = \dim_K \Delta(j)_k = \dim_K \left(P(j)_k / \left(\sum_{j < i} P(i) \right)_k \right) = [\Delta(j) : S(k)]$.

❷ Let $\mathbf{i} \in \mathcal{L}(j) := \{(i_1, i_2, \dots, i_r) \mid i_m \in \Lambda^{(j)}, i_l \not\leq i_t, 1 \leq l < t \leq r := |\Lambda^{(j)}|\}$ (see [9, 4.2]). Obviously, $P(i_t) \subseteq P(j) = P(i_1)$ for all $1 \leq t \leq r$. Denote by $\mathcal{D}(\mathbf{i})$ the filtration $0 = D(r+1) \subset D(r) \subset \dots \subset D(t) \subset \dots \subset D(1)$ with $D(t) := \sum_{m=t}^r P(i_m)$. It is easy to check that $\mathbf{B}_j(\Lambda^{(i_t)} \setminus \{i_t\}) = \mathbf{B}_j(\{i_t\}) \cap \mathbf{B}_j(\{i_{t+1}, \dots, i_r\})$ for all $1 \leq t \leq r-1$. This implies $\sum_{i_t < k} P(k) = P(i_t) \cap \left(\sum_{m=t+1}^r P(i_m) \right)$ (see 2.2.4(2)) and consequently $D(t)/D(t+1) \cong P(i_t) / \left(\sum_{i_t < k} P(k) \right) \cong \Delta(i_t)$ for all $1 \leq t \leq r$. The filtration $\mathcal{D}(\mathbf{i})$ of $P(j)$ is Δ -good. Since $\{i_1, i_2, \dots, i_r\} = \Lambda^{(j)}$ and $l \neq t$ implies $i_l \neq i_t$, we obtain $(P(j) : \Delta(i)) = 1$ for any $i \in \Lambda^{(j)}$ and $(P(j) : \Delta(i)) = 0$ if $i \in \Lambda \setminus \Lambda^{(j)}$.

❸ Since $P(i) \hookrightarrow P(1)$ for all $i \in \Lambda$, it is enough to show that $\text{soc } P(1) \cong S(1)$. We consider the map $f_{(1, n, 1)} : P(1) \rightarrow P(1)$ induced by $f_{(1, n, 1)} : L(1) \twoheadrightarrow L(n) \hookrightarrow L(1)$, here $L(n) = \text{soc } L(1)$ since n is maximal.

We show that $\text{im}(f_{(1, n, 1)}) \subseteq \text{im}(f)$ for all $f \in \text{Hom}_A(P(i), P(1))$ with $f \neq 0$ and all $i \in \Lambda$. This implies, that $\text{im}(f_{(1, n, 1)})$ is contained in every local submodule of $P(1)$ and therefore in every non-zero submodule of $P(1)$. Thus $\text{im}(f_{(1, n, 1)})$ is the uniquely determined simple submodule of $P(1)$ and $\text{top}(\text{im}(f_{(1, n, 1)})) \cong S(1)$ implies $\text{im}(f_{(1, n, 1)}) \cong S(1)$: Let $f \in \text{Hom}_A(P(i), P(1)) \setminus \{0\}$ be induced by $\mathbf{f} \in \text{Hom}_B(L(i), L(1))$, then $\mathbf{f} \neq 0$ and consequently $L(n) \subseteq \text{im}(\mathbf{f})$. There exists some $\mathbf{x} \in L(i)$ with $\mathbf{f}(\mathbf{x}) = \mathbf{b}_n$, where $\mathbf{b}_n$ is a generator of $L(n)$. Let $\mathbf{g}$ be in $\text{Hom}_B(L(1), L(i))$ given by $\mathbf{g} : L(1) \xrightarrow{\cdot \mathbf{x}} L(i)$ and $\mathbf{g} \in \text{Hom}_A(P(1), P(i))$ is induced by $\mathbf{g}$. We have $\mathbf{f} \circ \mathbf{g} = f_{(1, n, 1)}$. This implies $f \circ \mathbf{g} = f_{(1, n, 1)}$ and consequently $\text{im}(f_{(1, n, 1)}) \subseteq \text{im}(f)$.

According to 2.2.3 for the algebra $A^{op} \cong \text{End}_B(L) \cong \text{End}_{B^{op}}(\mathcal{D}(L))$ we have $\text{soc } P_{A^{op}}(i) \cong$

$S_{A^{op}}(1)$, thus $\text{top } I(i) \cong S(1)$ for all $i \in \Lambda$.

④ Let $\mathfrak{f}_{(n,n,j)} : P(j) \rightarrow P(n)$ be the A -map induced by the B -map $\mathfrak{f}_{(n,n,j)} : L(j) \twoheadrightarrow L(n)$. It is enough to show the equation $\sum_{j < i} P(i) = \ker(\mathfrak{f}_{(n,n,j)})$. This implies $P(j)/\left(\sum_{j < i} P(i)\right) \stackrel{\bullet}{=} \Delta(j) \hookrightarrow P(n) = \Delta(n)$ for any $j \in \Lambda$: Let $i \in \Lambda^{(j)} \setminus \{j\}$, then the functor $\text{Hom}_B(L, -)$ maps a B -map $\mathfrak{f} : \left(L(i) \xrightarrow{\mathfrak{f}_{(j,i,i)}} L(j) \xrightarrow{\mathfrak{f}_{(n,n,j)}} L(n)\right)$ to the A -map $\mathfrak{f} : \left(P(i) \xrightarrow{\mathfrak{f}_{(j,i,i)}} P(j) \xrightarrow{\mathfrak{f}_{(n,n,j)}} P(n)\right)$. Since $L(n) = \text{soc } B$ is simple, we have $\ker(\mathfrak{f}_{(n,n,j)}) = \text{rad } L(j)$, thus $L(i) \subseteq \text{rad } L(j)$ since $i \in \Lambda^{(j)} \setminus \{j\}$. Hence $\mathfrak{f}$ and therefore $\mathfrak{f}$ are zero-maps. Consequently, the submodule $P(i)$ of $P(j)$ belongs to $\ker(\mathfrak{f}_{(n,n,j)})$ for any $i \in \Lambda^{(j)} \setminus \{j\}$. We obtain $\sum_{j < i} P(i) \subseteq \ker(\mathfrak{f}_{(n,n,j)})$.

Because $\Delta(j) = P(j)/\left(\sum_{j < i} P(i)\right)$, there exists a submodule U of $\Delta(j)$ such that $P(j)/(\ker \mathfrak{f}_{(n,n,j)}) \cong \Delta(j)/U$. For the K -subspace of $\text{im}(\mathfrak{f}_{(n,n,j)})$ corresponding to some $k \in \Lambda$ we have $\dim_K(\text{im } \mathfrak{f}_{(n,n,j)})_k = \dim_K(P(j)/\ker \mathfrak{f}_{(n,n,j)})_k = \dim_K(\Delta(j)/U)_k \leq \dim_K(\Delta(j))_k$.

Let $k \in \Lambda_{(j)}$, then the B -map $\mathfrak{g} : \left(L(k) \xrightarrow{\mathfrak{f}_{(j,j,k)}} L(j) \xrightarrow{\mathfrak{f}_{(n,n,j)}} L(n)\right)$ is non-zero, thus the induced A -map $\mathfrak{g} : \left(P(k) \xrightarrow{\mathfrak{f}_{(j,j,k)}} P(j) \xrightarrow{\mathfrak{f}_{(n,n,j)}} P(n)\right)$ is non-zero. Hence $0 \neq \mathfrak{g}(e_k) \in (\text{im } \mathfrak{f}_{(n,n,j)})_k$, thus $\dim_K(\text{im } \mathfrak{f}_{(n,n,j)})_k \neq 0$ and consequently $(\text{im } \mathfrak{f}_{(n,n,j)})_k = (\Delta(j))_k$ for all $k \in \Lambda$, because $\dim_K(\Delta(j))_k \stackrel{\bullet}{=} \begin{cases} 1 & \text{if } k \in \Lambda_{(j)}, \\ 0 & \text{else} \end{cases}$. We obtain $U = 0$ and $\sum_{j < i} P(i) = \ker(\mathfrak{f}_{(n,n,j)})$.

Proposition 2.2.3 implies that the functor $\text{Hom}_B(L, -) : \text{mod } B \rightarrow \text{mod } A^{op}$ yields $\Delta_{A^{op}}(j) \hookrightarrow \Delta_{A^{op}}(n)$. By applying the duality we get $\nabla(n) \twoheadrightarrow \nabla(j)$ for all $j \in \Lambda$.

This finishes the proof of Theorem A. $\square$

The features of the B -module L , for a pair (B, L) with $\boxed{\leq}$, implies some properties for the quiver and relations of the corresponding 1-quasi-hereditary algebra $\text{End}_B(L)^{op}$.

2.2.5 Remark. Let (B, L) be a pair with the property $\boxed{\leq}$. For every $i \in \Lambda$ we fix a generator $x_i \in B$ of the direct summand $L(i)$ of L and $x_1 = 1 := 1_B$, here $\{1\} = \min(\Lambda, \leq)$. For all $i, j \in \Lambda$ with $j < i$ there exists $b_{ij} \in B$ with $x_j \cdot b_{ij} = x_i$, because $x_i \in L(i) \subset L(j) = B \cdot x_j = x_j \cdot B$ (see 2.2.1(1) and (3)).

We define an injective and a surjective B -map between $L(i)$ and $L(j)$ by

$$f_{(i \rightarrow j)} : L(i) \xrightarrow{\cdot 1} L(j) \quad \text{and} \quad f_{(i \leftarrow j)} : L(j) \xrightarrow{\cdot b_{ij}} L(i)$$

For any $l, t \in \Lambda$ let $\mathfrak{X}(L(l), L(t))$ be the space of non-invertible maps $f \in \text{Hom}_B(L(l), L(t))$ with the property "if $f = f_2 \circ f_1$ factors through add L , then either f_1 is a split monomorphism or f_2 is a split epimorphism". The number of arrows from l to t in the quiver of the algebra $A = \text{End}_B(L)^{op}$ is $\dim_K \mathfrak{X}(L(l), L(t))$ (see [1] or [2]).

According to 2.2.1(4), any map $f \in \text{Hom}_B(L(l), L(t))$ factors through $\bigoplus_{i \in \Lambda^{(l)} \cap \Lambda^{(t)}} L(i)$. If l and t are incomparable, then $l, t \notin \Lambda^{(l)} \cap \Lambda^{(t)}$, thus $\mathfrak{X}(L(l), L(t)) = 0$. Assume $l < t$ and $f : L(l) \rightarrow L(t)$, then there exists $b \in B$ with $x_t \cdot b = f(x_l)$, since $L(t) = B \cdot x_t \stackrel{2.2.1(1)}{=} x_t \cdot B$ and hence $f : \left(L(l) \twoheadrightarrow L(t) \xrightarrow{\cdot b} L(t)\right)$. If f is not surjective, then b is not invertible and consequently $f \notin \mathfrak{X}(L(l), L(t))$. If f is surjective but l and t are not the adjacent, then $f : L(l) \twoheadrightarrow L(j) \twoheadrightarrow L(t)$ for some $j \in \Lambda$ with $l < j < t$ and therefore $f \notin \mathfrak{X}(L(l), L(t))$.

Let $l \triangleleft t$ and $g : L(l) \rightarrow L(t)$ with $g : \left(L(l) \rightarrow L(t) \xrightarrow{\cdot b} L(t) \right)$, then $L(t) \xrightarrow{\cdot b} L(t)$ is a split epimorphism if $b = c \cdot 1$ for some $c \in K \setminus \{0\}$, in other words $g = c \cdot f_{(t \leftarrow l)}$. Using analogous arguments also for $j > t$, we obtain $\mathfrak{X}(L(l), L(t)) = \begin{cases} \text{span}_K \{f_{(l \rightarrow t)}\} & \text{if } l \triangleright t, \\ \text{span}_K \{f_{(t \leftarrow l)}\} & \text{if } l \triangleleft t, \\ 0 & \text{else.} \end{cases}$

(1) In the quiver $Q_0(A) = \Lambda$ of the 1-quasi-hereditary algebra A two vertices i and j are connected by an arrow if they are neighbours with respect to $\leq$, more precisely, we have $i \rightleftharpoons j$. Assume $j \triangleleft i$, then the B -maps $f_{(i \rightarrow j)}$ and $f_{(i \leftarrow j)}$ can be considered as the maps corresponding to the arrows $i \rightarrow j$ and $j \rightarrow i$ respectively. In this case we use the notation $L(i) \xrightleftharpoons[\cdot b_{ij}]{\cdot 1} L(j)$. In general the notation $L(l) \xrightleftharpoons[\cdot d]{\cdot b} L(t)$ means that l and t are neighbours and

$$(b, d) = \begin{cases} (1, b_{lt}) & \text{if } t \triangleleft l, \\ (b_{tl}, 1) & \text{if } t \triangleright l. \end{cases} \quad \text{We always have } (x_l \cdot b, x_t \cdot d) = \begin{cases} (x_l, x_l) & \text{if } t \triangleleft l, \\ (x_t, x_t) & \text{if } t \triangleright l. \end{cases}$$

(2) Let $p_t = (i, i_1^{(t)}, \dots, i_{m_t}^{(t)}, j)$ for $1 \leq t \leq r$ be some paths in $Q(A)$ (obviously, $i_k^{(t)}$ and $i_{k+1}^{(t)}$ are neighbours). Then

$$\sum_{t=1}^r c_t \cdot p_t \in \mathcal{I}(A) \quad \text{if and only if} \quad \sum_{t=1}^r c_t \cdot \left(L(i) \xrightarrow{\cdot b_0^{(t)}} L(i_1^{(t)}) \xrightarrow{\cdot b_1^{(t)}} \dots \xrightarrow{\cdot b_{m_t-1}^{(t)}} L(i_{m_t}^{(t)}) \xrightarrow{\cdot b_{m_t}^{(t)}} L(j) \right) = 0,$$

here the maps $L(i_k^{(t)}) \xrightarrow{\cdot b_k^{(t)}} L(i_{k+1}^{(t)})$ are of the form $f_{(l \rightarrow t)}$ or $f_{(l \leftarrow t)}$.

2.2.6 Lemma. *Let $A \cong \text{End}_B(L)^{op} \cong KQ/\mathcal{I}$ with $(\Lambda, \leq)$ be a 1-quasi-hereditary algebra, where $(B, L = \bigoplus_{i \in \Lambda} L(i))$ satisfies the property $\boxed{\leq}$.*

(1) *If p and q are some paths in Q of the form $p(j, i, k)$ (see Subsection 2.1), then $p - q \in \mathcal{I}$. In particular, $\mathfrak{o}(p(j, i, k)) = p(k, i, j)$.*

(2) *Let Γ be the set of (large) neighbours of 1, where $\{1\} = \min(\Lambda, \leq)$ and x_i be a generator of $L(i)$ for any $i \in \Gamma$. Then the set $\{x_i \mid i \in \Gamma\}$ is generating system of B . In particular, B is a factor algebra of $K\langle y_1, \dots, y_m \rangle$, where $m = |\Gamma|$.*

Proof. (1) Let $j < i$ and p, q be some increasing paths from j to i as well as $\mathfrak{o}(p), \mathfrak{o}(q)$ be the corresponding decreasing paths from j to i in Q , i.e., there exists $i = i_0 \triangleleft i_1 \triangleleft \dots \triangleleft i_m = j$ and $i = j_0 \triangleleft j_1 \triangleleft \dots \triangleleft j_r = j$ with $p = (i, i_1, \dots, i_m, j)$ and $q = (i, j_1, \dots, j_r, j)$ as well as $\mathfrak{o}(p) = (j, i_m, \dots, i_1, i)$ and $\mathfrak{o}(q) = (j, j_r, \dots, j_1, i)$. For the corresponding B -maps $f_{(p)} = f_{(i_m \rightarrow j)} \circ \dots \circ f_{(i_1 \rightarrow i_2)} \circ f_{(i \rightarrow i_1)}$ and $f_{(q)} = f_{(j_r \rightarrow j)} \circ \dots \circ f_{(j_1 \rightarrow j_2)} \circ f_{(i \rightarrow j_1)}$ as well as $f_{\mathfrak{o}(p)} = f_{(i \leftarrow i_1)} \circ f_{(i_1 \leftarrow i_2)} \circ \dots \circ f_{(i_m \leftarrow j)}$ and $f_{\mathfrak{o}(q)} = f_{(i \leftarrow j_1)} \circ f_{(j_1 \leftarrow j_2)} \circ \dots \circ f_{(j_r \leftarrow j)}$ we obtain $f_{(p)} - f_{(q)} = 0$ and $f_{\mathfrak{o}(p)} - f_{\mathfrak{o}(q)} = 0$.

For some $i, j, k \in \Lambda$ with $i \leq j, k$ let p and q be some paths in Q of the form $p(j, i, k)$, then $p = \mathfrak{o}(p_1) \cdot p_2$ and $q = \mathfrak{o}(q_1) \cdot q_2$ with some increasing paths p_1, q_1 from k to i and p_2, q_2 from j to i . For the corresponding B -maps we have $f_{(p)} = f_{(\mathfrak{o}(p_1))} \circ f_{(p_2)}$ and $f_{(q)} = f_{(\mathfrak{o}(q_1))} \circ f_{(q_2)}$. Since $f_{(p_1)} = f_{(q_1)}$ and $f_{(\mathfrak{o}(p_1))} = f_{(\mathfrak{o}(q_1))}$, we obtain $f_{(p)} = f_{(q)}$. This implies $p - q \in \mathcal{I}$.

(2) For any $i \in \Lambda$ with $i \neq 1$ there exists $j \in \Gamma$ with $j \leq i$. Thus $L(i) \subseteq L(j)$ and consequently $\text{rad } B = \sum_{i \in \Lambda \setminus \{1\}} L(i) = \sum_{i \in \Gamma} L(i)$ (see 1.2.4(2)(b)). The set $\{x_i \mid i \in \Gamma\}$ generates $\text{rad } B$. Since B is local, we obtain that B is a factor algebra of $K\langle y_1, \dots, y_{|\Gamma|} \rangle$. $\square$

3. Proof of Theorem B

In this section $A = KQ/\mathcal{I}$ with $(\Lambda, \leq)$ is a 1-quasi-hereditary algebra and $(B, L = \bigoplus_{i \in \Lambda} L(i))$ is the corresponding pair with the property $\boxed{\leq}$, i.e., $A \cong \text{End}_B(L)^{op}$. For a relation $\rho = \sum_{t=1}^r c_t \cdot p_t$ in $\mathcal{I}$ we define $\mathfrak{o}(\rho) = \sum_{t=1}^r c_t \cdot \mathfrak{o}(p_t)$. The definition of a $\text{BGG}(\Rightarrow)$ -algebra is given in Subsection 1.3.

For the proof of Theorem B we have to show the equivalence of the following statements:

- ❶ B is commutative, ❷ A is a $\text{BGG}(\Rightarrow)$ -algebra, ❸ $\rho \in \mathcal{I}$ if and only if $\mathfrak{o}(\rho) \in \mathcal{I}$

Let x_i be a generator of $L(i)$ for any $i \in \Lambda$ and $x_1 = 1 := 1_B$ where $\{1\} = \min(\Lambda, \leq)$. For $i, j \in \Lambda$ with $i \triangleleft j$ or $i \triangleright j$ we denote by $L(i) \xrightleftharpoons[\cdot d]{\cdot b} L(j)$ the B -maps described in 2.2.5.

3.1 Lemma. *The following statements are equivalent:*

(i) B is commutative.

(ii) Let $p = (i_0, i_1, \dots, i_m)$ be some path in Q and $L(i_0) \xrightleftharpoons[\cdot d_1]{\cdot b_1} L(i_1) \xrightleftharpoons[\cdot d_2]{\cdot b_2} \dots \xrightleftharpoons[\cdot d_m]{\cdot b_m} L(i_m)$ be the corresponding B -maps with $x_0 := x_{i_0}$ and $x_m := x_{i_m}$. Then

$$x_0 \cdot b_1 \cdot b_2 \cdots b_m = x_m \cdot d_m \cdots d_2 \cdot d_1$$

(iii) Statement (ii) holds for $m = 4$.

Proof. (i) $\Rightarrow$ (ii) We show this by induction on m : If $m = 1$, then for $L(i_0) \xrightleftharpoons[\cdot d_1]{\cdot b_1} L(i_1)$ we have $(x_0 \cdot b_1, x_1 \cdot d_1) \in \left\{ (x_0, x_0), (x_1, x_1) \right\}$ (see 2.2.5(1)), thus $x_0 \cdot b_1 = x_1 \cdot d_1$. Assume $x_0 \cdot b_1 \cdot b_2 \cdots b_{m-1} = x_{m-1} \cdot d_{m-1} \cdots d_2 \cdot d_1$, then by multiplication with b_m we obtain

$$x_0 \cdot b_1 \cdot b_2 \cdots b_{m-1} \cdot b_m = x_{m-1} \cdot b_m \cdot d_{m-1} \cdots d_2 \cdot d_1, \quad (\otimes)$$

because B is commutative. If $i_m \triangleleft i_{m-1}$, then $b_m = 1$ and $x_m \cdot d_m = x_{m-1}$. The equation $(\otimes)$ implies $x_0 \cdot b_1 \cdot b_2 \cdots b_{m-1} \cdot b_m = x_m \cdot d_m \cdot d_{m-1} \cdots d_2 \cdot d_1$. If $i_m \triangleright i_{m-1}$, we obtain $x_{m-1} \cdot b_m = x_m$ and $d_m = 1$, thus the equation $(\otimes)$ is $x_0 \cdot b_1 \cdot b_2 \cdots b_{m-1} \cdot b_m = x_m \cdot d_m \cdot d_{m-1} \cdots d_2 \cdot d_1$.

(ii) $\Rightarrow$ (iii) This is trivial.

(iii) $\Rightarrow$ (i) Let $i, j \in \{l \in \Lambda \mid 1 \triangleleft l\}$, then for $L(1) \xrightleftharpoons[\cdot d_1]{\cdot b_1} L(i) \xrightleftharpoons[\cdot d_2]{\cdot b_2} L(1) \xrightleftharpoons[\cdot d_3]{\cdot b_3} L(j) \xrightleftharpoons[\cdot d_4]{\cdot b_4} L(1)$, since $1 \triangleleft i \triangleright 1 \triangleleft j \triangleright 1$, we have $(b_1, d_1) = (x_i, 1)$, $(b_2, d_2) = (1, x_i)$, $(b_3, d_3) = (x_j, 1)$, $(b_4, d_4) = (1, x_j)$ (see 2.2.5(1)). By the assumption, we have $x_1 \cdot b_1 \cdot b_2 \cdot b_3 \cdot b_4 = x_1 \cdot d_4 \cdot d_3 \cdot d_2 \cdot d_1$, thus we obtain $x_i \cdot x_j = 1 \cdot x_i \cdot 1 \cdot x_j \cdot 1 = 1 \cdot x_j \cdot 1 \cdot x_i \cdot 1 = x_j \cdot x_i$. Thus B is commutative, because $\{x_i \mid 1 \triangleleft i\}$ is a generating system of B (see 2.2.6(2)). $\square$

Proof ❶ $\Rightarrow$ ❸ Let $\rho = \sum_{t=1}^r c_t \cdot \left(i_0^{(t)}, i_1^{(t)}, \dots, i_{m_t}^{(t)} \right)$ be a relation of A with $i = i_0^{(t)}$, $j = i_{m_t}^{(t)}$ and $L(i_{v-1}^{(t)}) \xrightleftharpoons[\cdot d_v^{(t)}]{\cdot b_v^{(t)}} L(i_v^{(t)})$ the corresponding B -maps for all $1 \leq t \leq r$. We obtain $\sum_{t=1}^r c_t \cdot \left(L(i) \xrightarrow{\cdot b_1^{(t)}} L(i_1^{(t)}) \xrightarrow{\cdot b_2^{(t)}} \dots \xrightarrow{\cdot b_{m_t}^{(t)}} L(j) \right) = 0$ (see 2.2.5(2)). Hence, $\sum_{t=1}^r c_t \cdot$


$(x_i \cdot b_1^{(t)} \cdots b_{m_t}^{(t)}) = 0$. According to Lemma 3.1 we have $\sum_{t=1}^r c_t \cdot (x_j \cdot d_{m_t}^{(t)} \cdots d_1^{(t)}) = 0$. This implies $\sum_{t=1}^r c_t \cdot \left(L(j) \xrightarrow{\cdot d_{m_t}^{(t)}} \cdots \xrightarrow{\cdot d_2^{(t)}} L(i_1^{(t)}) \xrightarrow{\cdot d_1^{(t)}} L(i) \right) = 0$ and consequently $\mathfrak{o}(\rho) = \sum_{t=1}^r c_t \cdot (i_{m_t}^{(t)}, \dots, i_1^{(t)}, i_0^{(t)}) \in \mathcal{I}$.

③ $\Rightarrow$ ① Let $\Gamma := \{k \in \Lambda \mid 1 \triangleleft k\}$. It is enough to show $x_k \cdot x_j = x_j \cdot x_k$ for all $k, j \in \Gamma$ (see 2.2.6): Let $k, j \in \Gamma$, then for the B -maps $f: (L(1) \xrightarrow{\cdot x_k} L(k) \xrightarrow{\cdot 1} L(1) \xrightarrow{\cdot x_j} L(j) \xrightarrow{\cdot 1} L(1))$ and $g: (L(1) \xrightarrow{\cdot x_j} L(j) \xrightarrow{\cdot 1} L(1) \xrightarrow{\cdot x_k} L(k) \xrightarrow{\cdot 1} L(1))$ we obtain $f = g$ if and only if $f(1) = x_k \cdot x_j = x_j \cdot x_k = g(1)$. Since f and g correspond to the paths $(1, k, 1, j, 1)$ and $(1, j, 1, k, 1)$ respectively, we have to show $(1, k, 1, j, 1) - (1, j, 1, k, 1) \in \mathcal{I}$ (or $(1, j, 1, k, 1) = (1, k, 1, j, 1)$ in A).

According to [9, Theorem 3.2] for the path $(j, 1, k)$ there exists some $c_i \in K$ with $\rho = (j, 1, k) - \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} c_i \cdot p(j, i, k) \in \mathcal{I}$. Since $\mathfrak{o}(j, 1, k) = (k, 1, j)$ and $\mathfrak{o}(p(j, i, k)) = p(k, i, j)$, by our assumption we obtain $\mathfrak{o}(\rho) = (k, 1, j) - \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} c_i \cdot p(k, i, j) \in \mathcal{I}$. Obviously, $(1, j, 1, k, 1) = (k \rightarrow 1) \cdot (j, 1, k) \cdot (1 \rightarrow j)$ and $(1, k, 1, j, 1) = \mathfrak{o}(1, j, 1, k, 1) = (j \rightarrow 1) \cdot (k, 1, j) \cdot (1 \rightarrow k)$. The relations ρ and $\mathfrak{o}(\rho)$ implies the following equations in A .

$$\begin{aligned} (1, j, 1, k, 1) &= \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} c_i \cdot ((k \rightarrow 1) \cdot p(j, i, k) \cdot (1 \rightarrow j)) \\ (1, k, 1, j, 1) &= \sum_{i \in \Lambda^{(j)} \cap \Lambda^{(k)}} c_i \cdot ((j \rightarrow 1) \cdot p(k, i, j) \cdot (1 \rightarrow k)) \end{aligned}$$


For every $i \in \Lambda^{(j)} \cap \Lambda^{(k)}$ the paths $p_{(i)} = (k \rightarrow 1) \cdot p(j, i, k) \cdot (1 \rightarrow j)$ and $q_{(i)} = (j \rightarrow 1) \cdot p(k, i, j) \cdot (1 \rightarrow k)$ are of the form $p(1, i, 1)$ (in the right picture the black and the gray path respectively). Thus Lemma 2.2.6(1) implies $p_{(i)} = q_{(j)}$ in A , hence $(1, j, 1, k, 1) = (1, k, 1, j, 1)$.



② $\Leftrightarrow$ ③ If the K -map $\mathfrak{o} : A \rightarrow A$ with $p \mapsto \mathfrak{o}(p)$ is an anti-automorphism of A and for some paths $p_1, \dots, p_r$, which start in i and end in j we have $\sum_{t=1}^r c_t \cdot p_t = 0$, then $\sum_{t=1}^r c_t \cdot \mathfrak{o}(p_t) = 0$ (in other words, if $\rho \in \mathcal{I}$, then $\mathfrak{o}(\rho) \in \mathcal{I}$).

On the other hand, if $\mathcal{I} = \{\mathfrak{o}(\rho) \mid \rho \in \mathcal{I}\}$, then $\mathfrak{o}$ is obviously an anti-automorphism. $\square$

3.2 Example. The pair $(B, L_{(i)})$ in the Example 1.2.6 provides a $\text{BGG}_{(\rightleftharpoons)}$ -algebra $A_i = \text{End}_B(L_i)^{op}$ for any $i = 1, 2, 3$, because B is commutative. The quiver and relations to the left and to the right present the algebra A_1 and A_2 respectively (the algebra A_3 is presented in [10, Example 1]). For any relation ρ of A_i also $\mathfrak{o}(\rho)$ is a relation.



4. Ringel-duality on generators-cogenerators of local self-injective algebras

Let $(\mathcal{A}, \leq)$ be a quasi-hereditary algebra, then for any $i \in Q_0(\mathcal{A})$ there exists a (up to isomorphism) uniquely determined indecomposable module $T(i) \in \mathfrak{F}(\Delta) \cap \mathfrak{F}(\nabla)$ with the following properties:

For all $j \in Q_0(\mathcal{A})$ with $j \not\leq i$ we have $[T(i) : S(j)] = 0$ and $[T(i) : S(i)] = (T(i) : \Delta(i)) = (T(i) : \nabla(i)) = 1$, moreover, $\mathfrak{F}(\Delta) \cap \mathfrak{F}(\nabla) = \text{add} \left(\bigoplus_{i \in Q_0(\mathcal{A})} T(i) \right)$, the module $T := \bigoplus_{i \in Q_0(\mathcal{A})} T(i)$ is called *characteristic tilting module*. The *Ringel-dual* $R(\mathcal{A}) := \text{End}_{\mathcal{A}}(T)^{op}$ of $\mathcal{A}$ is a basic algebra with $Q_0(R(\mathcal{A})) = Q_0(\mathcal{A})$ and quasi-hereditary with the opposite order $\geq$ (we use the notation $\leq_{(R)}$). Moreover, $R(R(\mathcal{A})) \cong \mathcal{A}$ as quasi-hereditary algebras (for more details, see [11]).

The functor $\mathcal{R} := \text{Hom}_{\mathcal{A}}(T, -) : \text{mod } \mathcal{A} \rightarrow \text{mod } R(\mathcal{A})$ yields an exact equivalence between the subcategories $\mathfrak{F}(\Delta_{\mathcal{A}})$ and $\mathfrak{F}(\nabla_{R(\mathcal{A})})$. Moreover, $P_{R(\mathcal{A})}(i) \cong \mathcal{R}(T(i))$, $T_{R(\mathcal{A})}(i) \cong \mathcal{R}(I(i))$ and $\Delta_{R(\mathcal{A})}(i) \cong \mathcal{R}(\nabla(i))$ for all $i \in Q_0(\mathcal{A})$.

The class of 1-quasi-hereditary algebras is not closed under Ringel duality. Example 4 in [10] presents a 1-quasi-hereditary algebra A for which $R(A)$ is not 1-quasi-hereditary. However, the properties of $R(A)$ for a 1-quasi-hereditary algebra A considered in [9, Lemma 6.2] yield the following lemma. For $R(A)$ -modules we will use the index (R) , (note that $1 \leq i \leq n$ implies $n \leq_{(R)} i \leq_{(R)} 1$ for all $i \in \Lambda := Q_0(R(A)) = Q_0(A)$).

4.1 Lemma. *Let $(A, \leq)$ be a 1-quasi-hereditary algebra with $1 \leq i \leq n$ for any $i \in \Lambda$ and let $(R(A), \leq_{(R)})$ be the Ringel dual of $(A, \leq)$. Then $\text{dom. dim } R(A) \geq 2$ and $P_{(R)}(n)$ is a minimal faithful $R(A)$ -module.*

Proof. Since $\{1\} = \max(\Lambda, \leq_{(R)})$, the definition of standard modules implies $P_{(R)}(1) = \Delta_{(R)}(1)$. According to [9, Lemma 6.2 (c) and (b)] we obtain $\text{soc}(\Delta_{(R)}(1)) \cong S_{(R)}(n)$ and $\Delta_{(R)}(i) \hookrightarrow \Delta_{(R)}(1)$, since $T(1) \cong S(1)$ (see [9, 5.3]). Consequently, for all $i \in \Lambda$ we have $\text{soc}(\Delta_{(R)}(i)) \in \text{add}(\text{soc } P_{(R)}(n))$, because $P_{(R)}(n) \cong I_{(R)}(n)$ (see [9, 6.2(a)]).

According to Remark 1.2.1, it is enough to show $P_{(R)}(i) \hookrightarrow P_{(R)}(n)^{r_i}$ (for some $r_i \in \mathbb{N}$) and $P_{(R)}(n)^{r_i}/P_{(R)}(i) \in \mathfrak{F}(\Delta_{(R)})$ for any $i \in \Lambda$: Since $T(i) \in \mathfrak{F}(\Delta)$, we have $\text{soc } T(i) \in \text{add} \left(\bigoplus_{j \in \Lambda} \text{soc } \Delta(j) \right) \stackrel{1.2.2}{=} \text{add}(S(1))$ for any $i \in \Lambda$. Let $\text{soc } T(i) \cong S(1)^{r_i}$, then $T(i) \hookrightarrow T(n)^{r_i}$ since $T(n) \cong I(1)$ (see [9, 5.3]). The exact sequence $\xi : 0 \rightarrow T(i) \rightarrow T(n)^{r_i} \rightarrow T(n)^{r_i}/T(i) \rightarrow 0$ yields $T(n)^{r_i}/T(i) \in \mathfrak{F}(\nabla)$, because $\mathfrak{F}(\nabla)$ is closed under cokernels of injective maps (see [11]). By applying $\mathcal{R}(-)$ to ξ we obtain an exact sequence $0 \rightarrow P_{(R)}(i) \rightarrow P_{(R)}(n)^{r_i} \rightarrow P_{(R)}(n)^{r_i}/P_{(R)}(i) \rightarrow 0$ with $P_{(R)}(n)^{r_i}/P_{(R)}(i) \in \mathfrak{F}(\Delta_{(R)})$ for any $i \in \Lambda$. $\square$


4.1. Transfer of Ringel duality

Throughout, we keep the notation of the sets $\mathbf{X}, \mathbf{Y}, \mathbf{X}(1), \mathbf{Y}(1)$ and the functions $\mathbf{X} \xrightleftharpoons[\Psi]{\Phi} \mathbf{Y}$ used in Section 1. Moreover, we denote by $\mathbf{X}(R(1))$ the set of isomorphism classes of Ringel-duals of 1-quasi-hereditary algebras. Lemma 4.1 implies that $\mathbf{X}(R(1)) \subseteq \mathbf{X}$. We denote by $\mathbf{Y}(R(1))$ the image of $\Phi|_{\mathbf{X}(R(1))}$. Moreover, let $\mathcal{X} := \mathbf{X}(1) \cup \mathbf{X}(R(1))$ and $\mathcal{Y} := \mathbf{Y}(1) \cup \mathbf{Y}(R(1))$ as well as $\widehat{\mathcal{X}} := \mathbf{X}(1) \cap \mathbf{X}(R(1))$ and $\widehat{\mathcal{Y}} := \mathbf{Y}(1) \cap \mathbf{Y}(R(1))$.

The map $\mathcal{R} : \mathcal{X} \rightarrow \mathcal{X}$ with $\mathcal{R}([A]) = [R(A)]$ is obviously bijective and $\mathcal{R}^2 = \text{id}_{\mathcal{X}}$. The Morita-Tachikawa Theorem 1.1 and Theorem A yield the transfer of Ringel-duality for $\mathcal{X}$ on $\mathcal{Y}$ (illustrated on the picture below).

Let $\mathbf{R} := \Phi|_{\mathcal{X}} \circ \mathcal{R}|_{\mathcal{X}} \circ \Psi|_{\mathcal{Y}}$, then

$$\begin{aligned} \mathbf{R} : \mathcal{Y} &\longrightarrow \mathcal{Y} \\ \mathbf{R}([\mathcal{B}, \mathcal{L}]) &:= [\tilde{R}(\mathcal{B}), \tilde{R}(\mathcal{L})] \\ \text{and} \\ \mathbf{R}(\mathbf{R}[\mathcal{B}, \mathcal{L}]) &= [\mathcal{B}, \mathcal{L}] \end{aligned}$$



Obviously, for any pair $[\mathcal{B}, \mathcal{L}] \in \mathcal{Y}$ the algebra $\mathcal{A} = \Psi([\mathcal{B}, \mathcal{L}])$ is 1-quasi-hereditary or $\mathcal{A}$ is the Ringel dual of a 1-quasi-hereditary algebra $R(\mathcal{A}) = \Psi([\tilde{R}[\mathcal{B}], \tilde{R}(\mathcal{L})])$. The minimal faithful $\mathcal{A}$ -module $\mathcal{L}$ is the projective indecomposable $P(1_{\mathcal{A}})$ which corresponds to the minimal vertex $1_{\mathcal{A}}$ (see 1.2.3 and 4.1). In particular, $1_{R(\mathcal{A})}$ is the maximal vertex in Λ with respect to the partial order corresponding to $\mathcal{A}$. The direct summands of the $\mathcal{B}$ -module $\mathcal{L}$ are isomorphic to $\text{Hom}_{\mathcal{A}}(P(i), P(1_{\mathcal{A}}))$ for any $i \in \Lambda$.

4.1.1 Lemma. *Let $[\mathcal{B}, \mathcal{L}] \in \mathcal{Y}$ and let $\mathcal{A}$ be a quasi-hereditary algebra with $[\mathcal{A}] = \Psi([\mathcal{B}, \mathcal{L}])$. Then for $\mathbf{R}([\mathcal{B}, \mathcal{L}]) := [\tilde{R}(\mathcal{B}), \tilde{R}(\mathcal{L})]$ the following hold:*

$$\mathcal{B} \cong \tilde{R}(\mathcal{B}) \quad \text{and} \quad \tilde{R}(\mathcal{L}) \cong \bigoplus_{i \in \Lambda} \text{Hom}_{\mathcal{A}}(T(i), T(1_{R(\mathcal{A})})).$$

In particular, there exists a 1-quasi-hereditary algebra A with $\mathcal{B} \cong \text{End}_A(P(1_A))^{op}$ and $\mathcal{L} \cong \bigoplus_{i \in \Lambda} \text{Hom}_A(P(i), P(1_A))$ or $\mathcal{L} \cong \bigoplus_{i \in \Lambda} \text{Hom}_A(T(i), T(n_A))$.

Proof. It follows from Theorem 1.1 (with the preceding notation) that $\mathcal{B} \cong \text{End}_{\mathcal{A}}(P(1_{\mathcal{A}}))^{op}$ and $\tilde{R}(\mathcal{B}) \cong \text{End}_{R(\mathcal{A})}(P(1_{R(\mathcal{A})}))^{op}$ as well as $\mathcal{L} \cong \bigoplus_{i \in \Lambda} \text{Hom}_{\mathcal{A}}(P(i), P(1_{\mathcal{A}}))$ and $\tilde{R}(\mathcal{L}) \cong \bigoplus_{i \in \Lambda} \text{Hom}_{R(\mathcal{A})}(P_{(R)}(i), P_{(R)}(1_{R(\mathcal{A})}))$. Since $R(\mathcal{A}) \cong \text{End}_{\mathcal{A}}(T)^{op}$ and $P_{(R)}(i) \cong \text{Hom}_{\mathcal{A}}(T, T(i))$, the functor $\text{Hom}_{\mathcal{A}}(T, -) : \text{mod } \mathcal{A} \rightarrow \text{mod } R(\mathcal{A})$ yields an isomorphism

$$\text{Hom}_{R(\mathcal{A})}(P_{(R)}(i), P_{(R)}(1_{R(\mathcal{A})})) \cong \text{Hom}_{\mathcal{A}}(T(i), T(1_{R(\mathcal{A})}))$$

for all $i \in \Lambda$ (see [1, 2.1]). In particular, $\mathcal{B} \cong \text{End}_{\mathcal{A}}(P(1_{\mathcal{A}}))^{op} \cong \text{End}_{R(\mathcal{A})}(P(1_{R(\mathcal{A})}))^{op} \cong \tilde{R}(\mathcal{B})$ because $P(1_{\mathcal{A}}) \cong T(1_{R(\mathcal{A})})$ (see [9, Remark 5.3 and Lemma 6.2 (a)]).

If the algebra $\mathcal{A}$ is 1-quasi-hereditary, then $\mathcal{L} \cong \bigoplus_{i \in \Lambda} \text{Hom}_{\mathcal{A}}(P(i), P(1_{\mathcal{A}}))$. If not, then $A = R(\mathcal{A})$ is 1-quasi-hereditary with $1_A \leq i \leq n_A$ for all $i \in Q_0(A)$. For $\Phi([A]) = [B, L]$ we obtain $[\mathcal{B}, \mathcal{L}] = \mathbf{R}([B, L]) = [B, \tilde{R}(L)]$. Therefore $\mathcal{B} \cong \text{End}_A(P(1_A))^{op}$ and $\mathcal{L} \cong \bigoplus_{i \in \Lambda} \text{Hom}_A(T(i), T(n_A))$. $\square$


Lemma 4.1.1 implies that for any $[\mathcal{B}, \mathcal{L}] \in \mathcal{Y}$ we have $\mathbf{R}([\mathcal{B}, \mathcal{L}]) = [\mathcal{B}, \tilde{R}(\mathcal{L})]$. In particular, $\tilde{R}(\mathcal{L})$ is a multiplicity-free generator-cogenerator of $\mathcal{B}$ (see 1.1). The Ringel-duality on $\mathcal{Y}$ conforms with the duality on a subclass of multiplicity-free generator-cogenerators of local self-injective algebras, which arises from 1-quasi-hereditary algebras (via $\mathcal{B} \cong \text{End}_A(P(1_A))^{op}$).

According to Theorem A the first component B of a pair in $\mathcal{Y}$ is a local self-injective algebra having a module satisfying the properties in Definition 1.2.4 for some partial order.


We denote by $\mathfrak{L}(B)$ the set of all B -modules M such that (B, M) or $(B, \tilde{R}(M))$ satisfies the condition $\boxed{\leq}$ for some partial order $\leq$ on $\Lambda = \{1, \dots, \dim_K B\}$. Obviously, there exist finitely many partial orders on Λ . However, there are examples of a partial order $\leq$ on Λ with infinitely many pairwise non-isomorphic B -modules M such that (B, M) satisfies the property $\boxed{\leq}$ (in the next example 4.1.2 the B -module L_2 depends on the choice of $\mu \in K$, there we have $\langle X^2 + \mu Y \rangle \not\cong \langle X^2 + \mu' Y \rangle$ if $\mu \neq \mu'$). Moreover, if (B, M) and (B, M') satisfy the property $\boxed{\leq}$, then $[B, M] \in \hat{\mathcal{Y}}$ does not imply $[B, M'] \in \hat{\mathcal{Y}}$.

Let $\mathbf{L}$ be the set of isomorphism classes of algebras B with $\mathfrak{L}(B) \neq \emptyset$ (the function $\mathbf{X}(1) \rightarrow \mathbf{L}$ with $[A] \mapsto [\text{End}_A(P(1))]$ is surjective, non-injective and the set $\mathbf{L}$ is not finite). For any $[B] \in \mathbf{L}$ we define $\mathcal{Y}(B) := \{[B, M] \mid M \in \mathfrak{L}(B)\}$ and $\mathcal{X}(B) := \{[\text{End}_B(M)^{op}] \mid M \in \mathfrak{L}(B)\}$. It is easy to see that for all $[B], [B'] \in \mathbf{L}$ with $[B] \neq [B']$ we have $\mathcal{Y}(B) \cap \mathcal{Y}(B') = \emptyset$ and a pair $[B, M] \in \mathcal{Y}$ belongs to $\mathcal{Y}(B)$. This implies $\mathcal{Y} = \bigcup_{[B] \in \mathbf{L}} \mathcal{Y}(B)$ and $\mathcal{X} = \bigcup_{[B] \in \mathbf{L}} \mathcal{X}(B)$.

In the picture the sets $\mathcal{X}$ and $\mathcal{Y}$ are presented as the disjoint union of $\mathcal{X}(B)$ and $\mathcal{Y}(B)$ (symbolized by the circles, they are closed under the Ringel-duality $\mathbf{R}$ and $\mathcal{R}$) respectively. The dark circles inside the circle corresponding to $\mathcal{X}(B)$ symbolizes $\mathcal{X}(B) \cap \hat{\mathcal{X}}$. Similarly, a pair $[B, M]$ in the dark circle of $\mathcal{Y}(B)$ has the property $\boxed{\leq}$ and $\mathbf{R}([B, M])$ has the property $\boxed{\geq}$. In particular, $\hat{\mathcal{X}}$ and $\hat{\mathcal{Y}}$ are the disjoint unions of the *dark circles*. They are also closed under $\mathcal{R}$ resp. $\mathbf{R}$. The observation of Ringel-duality on $\mathcal{X}$ and $\mathcal{Y}$ can be restricted to $\mathcal{X}(B)$ and $\mathcal{Y}(B)$ respectively. The dark circles inside the circle corresponding to $\mathcal{X}(B)$ symbolizes $\mathcal{X}(B) \cap \hat{\mathcal{X}}$. Similarly, a pair $[B, M]$ in the dark circle of $\mathcal{Y}(B)$ has the property $\boxed{\leq}$ and $\mathbf{R}([B, M])$ has the property $\boxed{\geq}$, i.e., $[B, M] \in \hat{\mathcal{Y}}$. In particular, $\hat{\mathcal{X}}$ and $\hat{\mathcal{Y}}$ are the disjoint unions of the *dark circles*. They are also closed under $\mathcal{R}$ resp. $\mathbf{R}$. The observation of Ringel-duality on $\mathcal{X}$ and $\mathcal{Y}$ can be restricted to $\mathcal{X}(B)$ and $\mathcal{Y}(B)$ respectively.



4.1.2 Example. The algebra $B = K[x, y] / \langle xy, x^4 - y^2 \rangle$ is local and self-injective with $\dim_K B = 6$. The pair (B, L_i) satisfies the property $\boxed{\leq_{(i)}}$, where $L_i = \bigoplus_{k=1}^6 L_i(k)$ and $\leq_{(i)}$ for $i = 1, 2$ are presented in the following diagrams in the same way as in Example 1.2.6. Both pairs belong to $\mathcal{Y}(B)$, however, $[B, L_1] \in \hat{\mathcal{Y}}$ and $[B, L_2] \notin \hat{\mathcal{Y}}$.



On the right-hand side we present the quiver and relations of the 1-quasi-hereditary algebra $A_1 = \text{End}_B(L_1)^{op}$. The algebras A_1 and $R(A_1)$ are isomorphic as quasi-hereditary algebras (i.e., A_1 is Ringel self-dual), because $[B, L_1] = [B, \tilde{R}(L_1)]$ (see Theorem 4.2.1). The quiver

and relations of $A_2 = \text{End}_B(L_2)^{op}$ can be found in [10, Example 4] (there $q = 1 + \mu^2$). The Ringel-dual $[B, \tilde{R}(L_2)]$ of $[B, L_2]$ is given by $\tilde{R}(L_2) = \bigoplus_{i=1}^6 \tilde{R}(L_2(i))$ with

$$\begin{aligned} \tilde{R}(L_2(1)) &= \langle X^4 \rangle, & \tilde{R}(L_2(2)) &= \langle X^3 \rangle, & \tilde{R}(L_2(3)) &= \langle \mu X^2 - Y \rangle, \\ \tilde{R}(L_2(4)) &= \langle Y \rangle, & \tilde{R}(L_2(5)) &= \langle (X, Y) \rangle + \langle (Y, 0) \rangle \subset B \oplus B, & \tilde{R}(L_2(6)) &= B. \end{aligned}$$

4.2. Ringel-duality on $\hat{\mathcal{Y}}$

In [9, Sec. 5 and 6] 1-quasi-hereditary algebras whose isomorphism classes belong to $\hat{\mathcal{X}}$ have been considered. These results imply a precise description of the Ringel-duality $\mathbf{R}$ on $\hat{\mathcal{Y}}$.

Let $[B, L]$ be in $\mathbf{Y}(1)$, then $B \cong \text{End}_A(P(1))^{op}$ and $L \cong \bigoplus_{i \in \Lambda} B \circ \mathbf{f}_{(i)}$, where $A = \text{End}_B(L)^{op}$ is a 1-quasi-hereditary algebra with $(\Lambda, \leq)$ (here $1 \leq i \leq n$ for all $i \in \Lambda$) and $\mathbf{f}_{(i)} := \mathbf{f}_{(1,i,1)}$ is the endomorphism of $P(1)$ corresponding to the path $p(1, i, 1)$ of A (see 2.1.1). Let $R(A) = \text{End}_A(T)^{op}$ with $(\Lambda, \leq_{(R)})$ also be 1-quasi-hereditary. According to Theorem 6.1 in [9], the direct summand $T(i)$ of the characteristic tilting A -module T is a submodule and a factor module of $P(1) \cong I(1)$ for any $i \in \Lambda$ or more precisely $T(i) \cong P(1) / \left(\sum_{l \in \Lambda \setminus \Lambda_{(i)}} P(l) \right) \cong \bigcap_{l \in \Lambda \setminus \Lambda_{(i)}} \ker(P(1) \twoheadrightarrow I(l))$. Consequently, the subspace of $P(1)$ corresponding to the vertex 1 contains an element $\mathbf{t}(i)$ which generates $T(i)$. For any $i \in \Lambda$ we denote by $\mathcal{T}_{(i)}$ the following endomorphism of $P(1)$:

$$\mathcal{T}_{(i)} = (\iota(i) \circ \pi(i)) : \left(P(1) \xrightarrow{\pi(i)} T(i) \xrightarrow{\iota(i)} P(1) \right) \text{ with } e_1 \xrightarrow{\pi(i)} \mathbf{t}(i) \text{ and } \mathbf{t}(i) \xrightarrow{\iota(i)} \mathbf{t}(i)$$

Using Lemma 3.2 [10] it is easy to show, that the pair $(B, \mathcal{L})$ with $\mathcal{L} \cong \bigoplus_{i \in \Lambda} B \circ \mathcal{T}_{(i)}$ satisfies the condition $\boxed{\leq_{(R)}}$. We recall that using the notations of Section 2, for all $i, j \in \Lambda$ with $i \not\leq j$ (and therefore $j \not\leq_{(R)} i$) we have $j \in \Lambda \setminus \Lambda^{(i)}$ resp. $i \in \Lambda \setminus \Lambda_{(j)}$.

Ringel-duality $\mathbf{R}$ on $\hat{\mathcal{Y}}$ provides a relationship between the endomorphisms $\mathcal{T}_{(1)}, \dots, \mathcal{T}_{(n)}$ and $\mathbf{f}_{(1)}, \dots, \mathbf{f}_{(n)}$ of $P(1)$. (The following statement yields Theorem C from the introduction.)

4.2.1 Theorem. *Let $(A, \leq)$ be a 1-quasi-hereditary algebra and let $[(B, L), \leq]$ be the corresponding pair in $\mathbf{Y}(1)$ with $B = \text{End}_A(P(1))^{op}$ and $L \cong \bigoplus_{i \in \Lambda} L(i)$ where $L(i) = B \circ \mathbf{f}_{(i)}$ for any $i \in \Lambda$. Let $\mathbf{R}([B, L]) = [B, \tilde{R}(L)]$ with $\tilde{R}(L) \cong \bigoplus_{i \in \Lambda} \tilde{R}(L(i))$. Then the following statements are equivalent:*

- (i) $[B, L] \in \hat{\mathcal{Y}}$.
- (ii) $\tilde{R}(L(i)) \cong B \circ \mathcal{T}_{(i)}$, where $\mathcal{T}_{(i)} \in B$ satisfies $\text{im } \mathcal{T}_{(i)} = T(i)$ for every $i \in \Lambda$.
- (iii) $\tilde{R}(L(i)) \cong B / \left(\sum_{j \in \Lambda \setminus \Lambda_{(i)}} L(j) \right) \cong \bigcap_{j \in \Lambda \setminus \Lambda_{(i)}} \ker(B \twoheadrightarrow L(j))$ for every $i \in \Lambda$.

Proof. Let A be a 1-quasi-hereditary algebra with $[A] = \Psi([B, L])$. According to 4.1.1 for any $i \in \Lambda$ we have $\tilde{R}(L(i)) \cong \text{Hom}_A(T(i), T(n))$.

(i) $\Rightarrow$ (ii) The assumption $[B, L] \in \hat{\mathcal{Y}}$ implies $[A] \in \hat{\mathcal{X}}$. As already explained, we can define endomorphisms $\mathcal{T}_{(i)} = (\iota(i) \circ \pi(i))$ for any $i \in \Lambda$. Since $\text{Hom}_A(-, P(1))$ is exact, the inclusion $T(i) \xrightarrow{\iota(i)} P(1)$ yields a surjective B -map $\text{Hom}_A(P(1), P(1)) \twoheadrightarrow \text{Hom}_A(T(i), P(1))$ with $F \mapsto F \circ \iota(i)$ and therefore $\text{Hom}_A(T(i), P(1)) = B \circ \iota(i)$ for all $i \in \Lambda$. The map $B \circ \iota(i) \rightarrow B \circ \mathcal{T}_{(i)}$ given by $F \circ \iota(i) \mapsto F \circ \iota(i) \circ \pi(i) = F \circ \mathcal{T}_{(i)}$ for all $F \in B$ is obviously a B -module isomorphism. We obtain $\tilde{R}(L(i)) \cong B \circ \iota(i) \cong B \circ \mathcal{T}_{(i)}$ for all $i \in \Lambda$.

(i) $\Leftarrow$ (ii) Since $\text{im } \mathcal{T}_{(i)} = T(i)$, the top of any direct summand of the characteristic tilting module of A is simple. According to Theorem 5.1 and 6.1 in [9] the algebra $(R(A), \leqslant_{(R)})$ is 1-quasi-hereditary. This implies $\mathbf{R}[B, L] \in \mathbf{Y}(1)$.

(i) $\Rightarrow$ (iii) Let $i \in \Lambda$ and $\Upsilon_{(i)} : B \twoheadrightarrow L(i)$ be the surjective B -map given by $\Upsilon_{(i)}(F) = F \circ \mathfrak{f}_{(i)}$ for all $F \in B$. Since $(\mathcal{T}_{(j)} \circ \mathfrak{f}_{(i)}) : \left(P(1) \xrightarrow{f_{(i,i,1)}} P(i) \xrightarrow{f_{(1,i,i)}} P(1) \xrightarrow{\pi(j)} T(j) \xrightarrow{\iota(j)} P(1) \right)$ (in the notations of Subsection 2.1) and since for any $j \in \Lambda \setminus \Lambda^{(i)}$ holds $\dim_K \text{Hom}_A(P(i), T(j)) = [T(j) : S(i)] = 0$ (because of the properties of $T(i)$), we obtain $\pi(j) \circ f_{(1,i,i)} = 0$ and hence $\mathcal{T}_{(j)} \circ \mathfrak{f}_{(i)} = 0$. For all $i, j \in \Lambda$ with $i \not\leqslant j$ we have $\tilde{R}(L(j)) = B \circ \mathcal{T}_{(j)} \subseteq \ker(\Upsilon_{(i)})$, thus

$$\sum_{j \in \Lambda \setminus \Lambda^{(i)}} \tilde{R}(L(j)) \subseteq \ker(\Upsilon_{(i)}) \quad \text{and} \quad \tilde{R}(L(j)) \subseteq \bigcap_{i \in \Lambda \setminus \Lambda_{(j)}} \ker(\Upsilon_{(i)})$$

By assumption the pairs (B, L) and $(B, \tilde{R}(L))$ satisfy the properties $\boxed{\leqslant}$ and $\boxed{\leqslant_{(R)}}$ respectively. Therefore $\dim_K L(i) \stackrel{2.2.1(1)}{=} |\Lambda^{(i)}|$ implies $\dim_K \ker(\Upsilon_{(i)}) = |\Lambda \setminus \Lambda^{(i)}|$, moreover, $\dim_K \left(\sum_{j \in \Lambda \setminus \Lambda^{(i)}} \tilde{R}(L(j)) \right) \stackrel{2.2.1(1)}{=} \left| \bigcup_{\substack{j \in \Lambda \\ j \not\leqslant_{(R)} i}} \{k \in \Lambda \mid j \leqslant_{(R)} k\} \right| = |\Lambda \setminus \Lambda^{(i)}|$. This implies $\sum_{j \in \Lambda \setminus \Lambda^{(i)}} \tilde{R}(L(j)) = \ker(\Upsilon_{(i)})$ and consequently $\bigcap_{i \in \Lambda \setminus \Lambda_{(j)}} \ker(\Upsilon_{(i)}) = \sum_{j \leqslant_{(R)} k} \tilde{R}(L(k)) = \tilde{R}(L(j))$, because $\tilde{R}(L(k)) \subseteq \tilde{R}(L(j))$ for all $k \in \Lambda$ with $j \leqslant_{(R)} k$. Moreover we obtain $L(i) \cong B / \ker(\Upsilon_{(i)}) \cong B / \left(\sum_{j \in \Lambda \setminus \Lambda^{(i)}} \tilde{R}(L(j)) \right) = B / \left(\sum_{j \not\leqslant_{(R)} i} \tilde{R}(L(j)) \right)$. Using the dual argumentation we obtain $\tilde{R}(L(i)) \cong B / \left(\sum_{j \in \Lambda \setminus \Lambda_{(i)}} L(j) \right)$ for any $i \in \Lambda$.

(iii) $\Rightarrow$ (i) We have to show that $(B, \tilde{R}(L) = \bigoplus_{i \in \Lambda} \tilde{R}(L(i)))$ satisfies the property $\boxed{\leqslant_{(R)}}$. Since $L(i) = B \circ \mathfrak{f}_{(i)} \stackrel{2.2.1(1)}{=} \mathfrak{f}_{(i)} \circ B$, for all $f, g \in B$ there exists $f' \in B$ with $g \circ f \circ \mathfrak{f}_{(i)} = g \circ \mathfrak{f}_{(i)} \circ f'$. Therefore, if $g \in \ker \left(B \xrightarrow{\text{of}_{(i)}} L(i) \right)$, then $g \circ f \in \ker \left(B \xrightarrow{\text{of}_{(i)}} L(i) \right)$ for all $f \in B$. The assumption $\tilde{R}(L(j)) \cong \bigcap_{i \in \Lambda \setminus \Lambda_{(j)}} \ker(B \twoheadrightarrow L(i))$ implies $\tilde{R}(L(j)) \circ B \subseteq \tilde{R}(L(j))$ and consequently $\tilde{R}(L(j))$ is two-sided local ideal of B , since $\tilde{R}(L(j))$ is a factor module and a submodule of B .

The property $\boxed{\leqslant}$ of $L(i)$ implies $\tilde{R}(L(k)) \twoheadrightarrow \tilde{R}(L(j))$ resp. $\tilde{R}(L(j)) \subseteq \tilde{R}(L(k))$ if and only if $k \leqslant_{(R)} j$, because $\Lambda \setminus \Lambda_{(k)} \subseteq \Lambda \setminus \Lambda_{(j)}$ if and only if $j \leqslant k$. We have $\text{rad } \tilde{R}(L(k)) = \sum_{k <_{(R)} j} \tilde{R}(L(j))$ for all $k \in \Lambda$ since $\tilde{R}(L(k)) / \left(\sum_{k <_{(R)} j} \tilde{R}(L(j)) \right)$ is simple. $\square$

4.2.2 Remark. If $[B, L] \in \hat{\mathcal{Y}}$, then the socle of $B / \left(\sum_{j \in \Lambda \setminus \Lambda_{(i)}} L(j) \right)$ is simple for all $i \in \Lambda$, because these factor modules of B are also submodules of B .

In Example 4.1.2 the pair $[B, L_2] \in \mathbf{Y}(1)$ is not in $\widehat{\mathcal{Y}}$, because for $i = 5$ we obtain $\text{soc} \left(B / \sum_{j \in \Lambda \setminus \Lambda_{(5)}} L_2(j) \right) = \text{soc} (B / L_2(6)) \cong (\text{soc}^2 B) / \text{soc} B \cong (\langle X^3 \rangle + \langle Y \rangle) / \langle X^4 \rangle \cong K \oplus K$.

4.2.3 Example. Let $B := B_n(C)$ be the algebra given in Example 1.3.2 and $L = B \oplus \bigoplus_{i=2}^{n-1} \langle X_i \rangle \oplus (\text{soc} B)$. The Ringel dual of the corresponding 1-quasi-hereditary algebra $A_n(C)$ is also 1-quasi-hereditary (see [10, Lemma 1.2]). Thus $[B, L] \in \widehat{\mathcal{Y}}$, in particular, $\widetilde{R}(L(1)) \cong \text{soc} B$, $\widetilde{R}(L(n)) \cong B$ and $\widetilde{R}(L(i)) \cong B / \left(\sum_{\substack{j=2 \\ i \neq j}}^{n-1} L(j) \right) \cong \left\langle \sum_{j=2}^{n-1} d_{ji} X_j \right\rangle$ for $2 \leq j \leq n-1$, where $C^{-1} = (d_{ij})_{2 \leq i, j \leq n-1}$.

4.2.4 Remark. If a 1-quasi-hereditary algebra A is Ringel self-dual, then $(A, \leq) \cong (R(A), \leq_{(R)})$ implies $[B, L] = \Phi([A]) = \Phi([R(A)]) = [B, \widetilde{R}(L)]$. In other words: The pair $[B, L]$ satisfies the property $\boxed{\leq}$ and the property $\boxed{\leq_{(R)}}$. Thus, there exists a permutation $\sigma \in \text{Sym}(\dim_K B)$ with $L(\sigma(i)) \cong \widetilde{R}(L(i))$.

The algebras associated with blocks of the category $\mathcal{O}(\mathfrak{g})$ are Ringel self-dual. In the Example 1.2.6 the 1-quasi-hereditary algebra $(A_3, \leq_{(3)})$ corresponding to the pair (B, L_3) is related to a regular block of $\mathcal{O}(\mathfrak{sl}_3)$. It is easy to check that the permutation $\sigma = (\sigma(1), \dots, \sigma(6)) = (6, 5, 4, 3, 2, 1) \in \text{Sym}(6)$ yields $L_3(\sigma(i)) \cong \widetilde{R}(L_3(i))$.

In the same example the algebra $(A_1, \leq_{(1)})$ is also Ringel self-dual. The permutation $\tau \in \text{Sym}(6)$ with $L_1(\tau(i)) \cong \widetilde{R}(L_1(i))$ is given by $\tau = (6, 4, 5, 2, 3, 1)$. The algebra $R(A_2)$ is not 1-quasi-hereditary, because $\text{soc}(B / \left(\sum_{j \in \Lambda \setminus \Lambda_{(4)}} L_2(j) \right)) \cong \text{soc} (B / L_2(5))$ is not simple.

In general, for some $[B, L] \in \widehat{\mathcal{Y}}$ the equation $[B, L] = [B, \widetilde{R}(L)]$ is not satisfied (see Example 4.2.3). Consequently a 1-quasi-hereditary algebra A with $[A] \in \widehat{\mathcal{X}}$ is not Ringel self-dual, in general.

Acknowledgments. I would like to thank Jean-Marie Bois, Rolf Farnsteiner, Julian Külshammer and Julia Worch for their helpful and constructive comments.

References

- [1] M. Auslander, I. Reiten, S. Smalø, *Representation Theory of Artin Algebras*, Cambridge Studies in Advanced Math. 36 (1995).
- [2] I. Assem, D. Simson, A. Skowronski, *Elements of the Representation Theory of Associative Algebras 1. Techniques of Representation Theory*, London Mathematical Society Student Texts 65, Cambridge University (2006).
- [3] E. Cline, B. Parshall and L. Scott, *Finite dimensional algebras and highest weight categories*, J. Reine Angew. Math. 391, 85-99 (1988). Math., 82, Amer. Math. Soc. (1989).
- [4] V. Dlab, P. Heath and F. Marko, *Quasi-hereditary endomorphism algebras*, Canad. Math. Bull. 38, 421-428 (1995).
- [5] V. Dlab and C.M. Ringel, *The module theoretic approach to quasi-hereditary algebras*, Representations of algebras and related topics (Kyoto, 1990), 200-224, London Math. Soc. Lecture Note Ser. 168, Cambridge Univ. Press, Cambridge (1992).
- [6] R. Irving, *BGG algebras and the BGG reciprocity principle*, J. Algebra 135, 363-380 (1990).

- [7] S. König, I.H. Slungård and C.C.Xi *Double centralizer properties, dominant dimension and tilting modules*, J. Algebra 240, 393-412 (2001).
- [8] M. Fang and S. König, *Schur functors and dominant dimension*, Trans. Amer. Math. Soc. 363, 1555-1576 (2011).
- [9] D. Pučinskaitė, *1-quasi-hereditary algebras*, J. London Math. Soc. (to appear).
- [10] D. Pučinskaitė, *1-quasi-hereditary algebras: Examples and invariant submodules of projectives*, <http://arxiv.org/pdf/1201.4311v1.pdf>
- [11] C.M. Ringel, *The category of modules with good filtrations over a quasi-hereditary algebra has almost split sequences*, Math. Z. 208, 209-223 (1991).
- [12] C.M. Ringel, *Artin algebras of dominant dimension at least 2*, <http://www.mathematik.uni-bielefeld.de/~ringel/opus/domdim.pdf>
- [13] C. Stroppel, *Category $\mathcal{O}$: Quivers and Endomorphism Rings of Projectives*, Represent. Theory 7, 322-345 (2003).
- [14] C. Xi, *Quasi-Hereditary algebras with a duality*, J. Reine angew. Math. 449, 201-205 (1994).